

Cochrane PAIN, PALLIATIVE & SUPPORTIVE CARE Reviews used to inform clinical guidelines published between 1st January 2018 to 31st March 2020

Name and date of clinical guideline	Title of review	DOI of version cited
Royal Osteoporosis Society. Vitamin D and Bone Health: A practical clinical guideline for patient management, Version 2. Bath: Royal Osteoporosis Society; 2020. [Issued December 2018; updated February 2020]. Available from: https://strwebprdmedia.blob.core.windows.net/media/ef2ideu2/ros-vitamin-d-and-bone-health-in-adults-february-2020.pdf	Vitamin D for the treatment of chronic painful conditions in adults	10.1002/14651858.CD007771.pub3
Ettinger DS, Berger MJ, Barbour S, Bergsbaken J, Brandt D, Brown GE, Crews JR, Kim EJ, Kirkegaard S, Kloth DD, Klute K, Kris M, Lim D, Loprinzi C, Ma CX, Maurer V, McAlister RK, Mehta R, Michaud LB, Navari RM, Noonan K, Riebandt G, Roeland E, Rugo HS, Saunders I, Scullion B, Silveira M, Wolf A Hughes M, Shead DA; National Comprehensive Cancer Network. NCCN Clinical Practice Guidelines in Oncology: Antiemesis, Version 1.2020. Plymouth (PA): National Comprehensive Cancer Network; 2020 February. Available from: https://www.nccn.org/professionals/physician_gls/pdf/antiemesis.pdf	Serotonin receptor antagonists for highly emetogenic chemotherapy in adults	10.1002/14651858.CD006272.pub2
Berger AM, Mooney K, Banerjee C, Breitbart WS, Carpenter KM, Chang Y, Cleeland C, Davis E, Dest V, DuBenske LL, Escalante CP, Fernandez-Robles C, Garcia S, Jankowski C, Jatoi A, Kinczewski LE, Loggers ET, Mandrell B, McInnes S, Meyer F, Murphy BA, Palesh O, Patel H, Riba MB, Roshal A, Rugo HS, Salvador C, Wagner-Johnston N, Walter M, Webb JA; National Comprehensive Cancer Network. NCCN Clinical Practice Guidelines in Oncology: Cancer-Related Fatigue, Version 1.2020. Plymouth (PA): National Comprehensive Cancer Network; 2020 February. Available from: https://www.nccn.org/professionals/physician_gls/pdf/fatigue.pdf	Exercise for the management of cancer-related fatigue in adults	10.1002/14651858.CD006145.pub2
	Drug therapy for the management of cancer-related fatigue	10.1002/14651858.CD006704.pub3
	Pharmacological treatments for fatigue associated with palliative care	10.1002/14651858.CD006788.pub2
	Psychosocial interventions for reducing fatigue during cancer treatment in adults	10.1002/14651858.CD006953.pub2
	Educational interventions for the management of cancer-related fatigue in adults	10.1002/14651858.CD008144.pub2
	Psychosocial interventions for fatigue during cancer treatment with palliative intent	10.1002/14651858.CD012030.pub2

<p>Riba MB, Donovan KA, Andersen B, Braun I, Breitbart WS, Brewer BW, Buchmann LO, Collins M, Corbett C, Fleishman S, Garcia S, Greenberg DB, Handzo GF, Hoofring L, Huang C-H, Kumar P, Lally R, Martin S, McGuffrey L, Mitchell W, Morrison LJ, Niazi SK, Pailler M, Palesh O, Parnes F, Pazar JP, Ralston L, Salman J, Shannon-Dudley MM, Valentine AD, Vanderlan J; National Comprehensive Cancer Network. NCCN Clinical Practice Guidelines in Oncology: Distress Management, Version 1.2020. Plymouth (PA): National Comprehensive Cancer Network; 2020 January. Available from: https://www.nccn.org/professionals/physician_gls/pdf/distress.pdf</p>	<p>Drug therapy for the management of cancer-related fatigue</p>	<p>10.1002/14651858.CD006704.pub3</p>
<p>Dans M, Kutner JS, Baker JN, Bauman JR, Beck AC, Campbell TC, Carey EC, Case AA, Dalal, Doberman DJ, Epstein AS, Fecher L, Jones J, Kapo J, Lee RT, Loggers ET, McCammon S, Miller C, Misra S, Mitchell W, Ogunseit AB, Portman DG, Sutton L, Temel J, Teply ML, Thomas J, Zachariah; National Comprehensive Cancer Network. NCCN Clinical Practice Guidelines in Oncology: Palliative Care, Version 1.2020. Plymouth (PA): National Comprehensive Cancer Network; 2020 February. Available from: https://www.nccn.org/professionals/physician_gls/pdf/palliative.pdf</p>	<p>Corticosteroids for the resolution of malignant bowel obstruction in advanced gynaecological and gastrointestinal cancer</p>	<p>10.1002/14651858.CD001219</p>
	<p>Surgery for the resolution of symptoms in malignant bowel obstruction in advanced gynaecological and gastrointestinal cancer</p>	<p>10.1002/14651858.CD002764.pub2</p>
	<p>Laxatives for the management of constipation in people receiving palliative care</p>	<p>10.1002/14651858.CD003448.pub4</p>
	<p>Megestrol acetate for treatment of anorexia-cachexia syndrome</p>	<p>10.1002/14651858.CD004310.pub3</p>
	<p>Haloperidol for the treatment of nausea and vomiting in palliative care patients</p>	<p>10.1002/14651858.CD006271.pub3</p>
	<p>Mu-opioid antagonists for opioid-induced bowel dysfunction in people with cancer and people receiving palliative care</p>	<p>10.1002/14651858.CD006332.pub3</p>
	<p>Pharmacological treatments for fatigue associated with palliative care</p>	<p>10.1002/14651858.CD006788.pub3</p>
	<p>Benzodiazepines for the relief of breathlessness in advanced malignant and non-malignant diseases in adults</p>	<p>10.1002/14651858.CD007354.pub3</p>
	<p>Effectiveness and cost-effectiveness of home palliative care services for adults with advanced illness and their caregivers</p>	<p>10.1002/14651858.CD007760.pub2</p>
	<p>Palliative pharmacological sedation for terminally ill adults</p>	<p>10.1002/14651858.CD010206.pub2</p>

	Opioids for the palliation of refractory breathlessness in adults with advanced disease and terminal illness	10.1002/14651858.CD011008.pub2
	Early palliative care for adults with advanced cancer	10.1002/14651858.CD011129.pub2
	Corticosteroids for adult patients with advanced cancer who have nausea and vomiting (not related to chemotherapy, radiotherapy, or surgery)	10.1002/14651858.CD012002.pub2
	Psychosocial interventions for fatigue during cancer treatment with palliative intent	10.1002/14651858.CD012030.pub2
AIM Specialty Health. Clinical Appropriateness Guidelines: Interventional Pain Management. Chicago, IL: AIM Specialty Health; 2020. [Effective January 2020; last reviewed September 2018]. Available from: https://aimspecialtyhealth.com/wp-content/uploads/2019/11/AIM_Guidelines_MSK_Interventional-Pain-Management-1.pdf	Local anaesthetic sympathetic blockade for complex regional pain syndrome	10.1002/14651858.CD004598.pub4
McCusker M, Jolkvosky M, Ruff R, Benson J, Rosielle DA, Schmidt M, Hansen A, Dvorkin J, Hadzic S. Health Care Guideline: Palliative Care for Adults (Sixth Edition). Bloomington, MN: Institute for Clinical Systems Improvement (ICSI); 2020. [Issued January 2020]. Available from: https://www.icsi.org/wp-content/uploads/2020/01/PalliativeCare_6th-Ed_2020_v2.pdf	Opioids for the palliation of refractory breathlessness in adults with advanced disease and terminal illness	10.1002/14651858.CD011008.pub2
Suomalaisen Lääkäriseuran Duodecimin, Suomen Psykiatriyhdistys ry:n asettama työryhmä, Finnish Medical Society Duodecim, Finnish Psychiatric Association. Duodecim Käypä Hoito: Depressio. [Duodecim Current Care Guidelines: Depression]. Helsinki: Finnish Medical Society Duodecim; January 2020. Available from: https://www.kaypahoito.fi/hoi50023	Antidepressants for neuropathic pain	10.1002/14651858.CD005454
Bremerich D, Annecke T, Chappell D, Hanß R, Kaufner L, Kehl F, Kranke P, Girard T, Gogarten W, Greve S, Neuhaus S, Schlembach D, Schlösser L, Standl T, Treskatsch S, Volk T, Wallenborn J, Weber S, Wenk M; Deutsche Gesellschaft für Anästhesiologie und Intensivmedizin e.V. (DGAI); Deutsche Gesellschaft für Gynäkologie	Drug therapy for preventing post-dural puncture headache	10.1002/14651858.CD001792.pub3
	Patient controlled opioid analgesia versus non-patient controlled opioid analgesia for postoperative pain	10.1002/14651858.CD003348.pub3
	Single dose oral celecoxib for acute postoperative pain in adults	10.1002/14651858.CD004233.pub4

<p>und Geburtshilfe e.V. (DGGG). S1-Leitlinie: Die geburtshilfliche Analgesie und Anästhesie. [S1-Guideline: Obstetric analgesia and anaesthesia]. Nürnberg: Deutsche Gesellschaft für Anästhesiologie und Intensivmedizin; 2020. [Issued March 2020]. Available from: https://www.awmf.org/leitlinien/detail/II/001-038.html</p>	<p>Drug therapy for treating post-dural puncture headache</p>	<p>10.1002/14651858.CD007887.pub3</p>
<p>Deutsche Schmerzgesellschaft e.V. S3-Leitlinie: Langzeitanwendung von Opioiden bei chronischen nicht-tumorbedingten Schmerzen (LONTS). [S3- guideline: Long-term use of opioids in chronic non-cancer-related pain]. Berlin: Deutsche Schmerzgesellschaft e.V.; 2020. Available from: https://www.awmf.org/leitlinien/detail/II/145-003.html</p>	<p>Tapentadol for chronic musculoskeletal pain in adults</p>	<p>10.1002/14651858.CD009923.pub2</p>
	<p>Transdermal fentanyl for cancer pain</p>	<p>10.1002/14651858.CD010270.pub2</p>
	<p>Interventions for the reduction of prescribed opioid use in chronic non-cancer pain</p>	<p>10.1002/14651858.CD010323.pub3</p>
	<p>Adverse events associated with medium- and long-term use of opioids for chronic non-cancer pain: an overview of Cochrane Reviews</p>	<p>10.1002/14651858.CD012509.pub2</p>
<p>Scottish Intercollegiate Guidelines Network (SIGN). Management of chronic pain. Edinburgh: SIGN: 2019. (Sign publication no. 136). [Issued December 2013; updated August 2019]. Available from: https://www.sign.ac.uk/assets/sign136_2019.pdf</p>	<p>Transcutaneous electrical nerve stimulation (TENS) for chronic pain</p>	<p>10.1002/14651858.CD003222.pub2</p>
	<p>Music for pain relief</p>	<p>10.1002/14651858.CD004843.pub2</p>
	<p>Topical lidocaine for the treatment of postherpetic neuralgia</p>	<p>10.1002/14651858.CD004846.pub2</p>
	<p>Carbamazepine for acute and chronic pain in adults</p>	<p>10.1002/14651858.CD005451.pub2</p>
	<p>Lamotrigine for acute and chronic pain</p>	<p>10.1002/14651858.CD006044.pub3</p>
	<p>Long-term opioid management for chronic noncancer pain</p>	<p>10.1002/14651858.CD006605.pub2</p>
	<p>Pregabalin for acute and chronic pain in adults</p>	<p>10.1002/14651858.CD007076.pub2</p>
	<p>Topical capsaicin (high concentration) for chronic neuropathic pain in adults</p>	<p>10.1002/14651858.CD007393.pub3</p>
	<p>Topical NSAIDs for chronic musculoskeletal pain in adults</p>	<p>10.1002/14651858.CD007400.pub2</p>
	<p>Topical rubefaciants for acute and chronic pain in adults</p>	<p>10.1002/14651858.CD007403.pub2</p>
	<p>Vitamin D for the treatment of chronic painful conditions in adults</p>	<p>10.1002/14651858.CD007771.pub2</p>
	<p>Gabapentin for chronic neuropathic pain and fibromyalgia in adults</p>	<p>10.1002/14651858.CD007938.pub2</p>
	<p>Amitriptyline for neuropathic pain and fibromyalgia in adults</p>	<p>10.1002/14651858.CD008242.pub2</p>
	<p>Milnacipran for neuropathic pain and fibromyalgia in adults</p>	<p>10.1002/14651858.CD008244.pub2</p>
	<p>Combination pharmacotherapy for the treatment of neuropathic pain in adults</p>	<p>10.1002/14651858.CD008943.pub2</p>

	Valproic acid and sodium valproate for neuropathic pain and fibromyalgia in adults	10.1002/14651858.CD009183.pub2
	Phenytoin for neuropathic pain and fibromyalgia in adults	10.1002/14651858.CD009485.pub2
	Clonazepam for neuropathic pain and fibromyalgia in adults	10.1002/14651858.CD009486.pub2
	Tapentadol for chronic musculoskeletal pain in adults	10.1002/14651858.CD009923.pub2
	Topical capsaicin (low concentration) for chronic neuropathic pain in adults	10.1002/14651858.CD010111
	Oxycodone for neuropathic pain in adults	10.1002/14651858.CD010692.pub3
	Buprenorphine for neuropathic pain in adults	10.1002/14651858.CD011603.pub2
	Hydromorphone for neuropathic pain in adults	10.1002/14651858.CD011604.pub2
	Morphine for chronic neuropathic pain in adults	10.1002/14651858.CD011669.pub2
	High-dose opioids for chronic non-cancer pain: an overview of Cochrane Reviews	10.1002/14651858.CD012299.pub2
	Methadone for neuropathic pain in adults	10.1002/14651858.CD012499.pub2
National Collaborating Centre for Cancer; National Institute for Health and Care Excellence (commissioner). Prostate cancer: Diagnosis and treatment. London: National Institute for Health and Care Excellence; 2019. (NICE NG131). [originally NICE CG175 issued January 2014; replaced by NG131 issued May 2019]. Available from: https://www.nice.org.uk/guidance/ng131	Radiotherapy for the palliation of painful bone metastases	10.1002/14651858.CD001793
	Radioisotopes for metastatic bone pain	10.1002/14651858.CD003347
	Interventions for sexual dysfunction following treatments for cancer	10.1002/14651858.CD005540.pub2
National Clinical Guideline Centre; National Institute for Health and Clinical Excellence (commissioner). Motor neurone disease: assessment and management. London: National Institute for Health and Clinical Excellence; 2019. (NICE NG42). [Issued February 2016; last updated July 2019]. Available from: https://www.nice.org.uk/guidance/ng42	Oxygen therapy for dyspnoea in adults	10.1002/14651858.CD004769.pub3
	Medically assisted nutrition for adult palliative care patients	10.1002/14651858.CD006274.pub3
	Benzodiazepines for the relief of breathlessness in advanced malignant and non-malignant diseases in adults	10.1002/14651858.CD007354.pub2
	Interventions for fatigue and weight loss in adults with advanced progressive illness	10.1002/14651858.CD008427.pub2
	Opioids for the palliation of refractory breathlessness in adults with advanced disease and terminal illness	10.1002/14651858.CD011008
National Guideline Alliance (hosted by the Royal College of Obstetricians and Gynaecologists); National Institute for Health and Care Excellence (commissioner). End of life care for infants, children and young people with life-limiting conditions: planning	Radiotherapy for the palliation of painful bone metastases	10.1002/14651858.CD001793
	Aromatherapy and massage for symptom relief in patients with cancer	10.1002/14651858.CD002287.pub2
	Oral morphine for cancer pain	10.1002/14651858.CD003868.pub4

and management. London: National Institute for Health and Care Excellence; 2019. (NICE NG61). [Issued December 2016; last updated July 2019]. Available from: https://www.nice.org.uk/guidance/ng61/	Local anaesthetic sympathetic blockade for complex regional pain syndrome	10.1002/14651858.CD004598.pub3
	Medically assisted nutrition for adult palliative care patients	10.1002/14651858.CD006274.pub3
	Music therapy for end-of-life care	10.1002/14651858.CD007169.pub2
	Benzodiazepines for the relief of breathlessness in advanced malignant and non-malignant diseases in adults	10.1002/14651858.CD007354.pub2
	Spiritual and religious interventions for well-being of adults in the terminal phase of disease	10.1002/14651858.CD007544.pub2
	Buprenorphine for treating cancer pain	10.1002/14651858.CD009596.pub4
	Psychological interventions for parents of children and adolescents with chronic illness	10.1002/14651858.CD009660.pub3
	Pharmacological interventions for pain in children and adolescents with life-limiting conditions	10.1002/14651858.CD010750.pub2
	Opioids for the palliation of refractory breathlessness in adults with advanced disease and terminal illness	10.1002/14651858.CD011008.pub2
National Clinical Guideline Centre; National Institute for Health and Care Excellence (commissioner). Chronic obstructive pulmonary disease in over 16s: diagnosis and management. London: Royal College of Physicians of London; 2019. (NICE NG115). [issued December 2018; last updated July 2019]. Available from: https://www.nice.org.uk/guidance/ng115	Opioids for the palliation of breathlessness in terminal illness	10.1002/14651858.CD002066
National Institute for Health and Care Excellence. Delirium: prevention, diagnosis and management. London: National Institute for Health and Care Excellence; 2019. (NICE CG103).[Issued July 2010; last updated March 2019]. Available from: https://www.nice.org.uk/guidance/cg103	Drug therapy for delirium in terminally ill adult patients	10.1002/14651858.CD004770.pub2
National Guideline Centre (hosted by the Royal College of Physicians); National Institute for Health and Care Excellence (commissioner). End of life care for adults: service delivery. London: National Institute for Health and Care Excellence; 2019. (NICE NG142). [Issued October 2019]. Available from: https://www.nice.org.uk/guidance/ng142	Interventions for improving palliative care for older people living in nursing care homes	10.1002/14651858.CD007132.pub2
	Effectiveness and cost-effectiveness of home palliative care services for adults with advanced illness and their caregivers	10.1002/14651858.CD007760.pub2
	End-of-life care pathways for improving outcomes in caring for the dying	10.1002/14651858.CD008006.pub4
	Early palliative care for adults with advanced cancer	10.1002/14651858.CD011129.pub2

National Institute for Health and Care Excellence. Clinical Knowledge Summaries: Headache - assessment. London: National Institute for Health and Care Excellence; 2019. [Last revised October 2019]. Available from: https://cks.nice.org.uk/headache-assessment	Ibuprofen for acute treatment of episodic tension-type headache in adults	10.1002/14651858.CD011474.pub2
	Paracetamol (acetaminophen) for acute treatment of episodic tension-type headache in adults	10.1002/14651858.CD011889.pub2
National Institute for Health and Care Excellence. Clinical Knowledge Summaries: Migraine. London: National Institute for Health and Care Excellence; 2019. [last revised April 2019]. Available from: http://cks.nice.org.uk/migraine	Acupuncture for the prevention of episodic migraine	10.1002/14651858.CD001218.pub3
	Zolmitriptan for acute migraine attacks in adults	10.1002/14651858.CD008616.pub2
National Institute for Health and Care Excellence. Clinical Knowledge Summaries: Chronic obstructive pulmonary disease. London: National Institute for Health and Care Excellence; 2019. [last revised August 2019]. Available from: http://cks.nice.org.uk/chronic-obstructive-pulmonary-disease	Benzodiazepines for the relief of breathlessness in advanced malignant and non-malignant diseases in adults	10.1002/14651858.CD007354.pub2
World Health Organization. WHO guidelines for the pharmacological and radiotherapeutic management of cancer pain in adults and adolescents. Geneva: World Health Organization; 2019. Available from: https://apps.who.int/iris/bitstream/handle/10665/279700/9789241550390-eng.pdf?ua=1	Bisphosphonates for the relief of pain secondary to bone metastases	10.1002/14651858.CD002068
	Oral morphine for cancer pain	10.1002/14651858.CD003868.pub4
	Oxycodone for cancer-related pain	10.1002/14651858.CD003870.pub5
	Methadone for cancer pain	10.1002/14651858.CD003971.pub3
	Codeine, alone and with paracetamol (acetaminophen), for cancer pain	10.1002/14651858.CD006601.pub4
	Buprenorphine for treating cancer pain	10.1002/14651858.CD009596.pub4
	Transdermal fentanyl for cancer pain	10.1002/14651858.CD010270.pub2
	Corticosteroids for the management of cancer-related pain in adults	10.1002/14651858.CD010756.pub2
	Hydromorphone for cancer pain	10.1002/14651858.CD011108.pub2
	Oral tapentadol for cancer pain	10.1002/14651858.CD011460.pub2
Wright M, Southcott E, MacLaughlin H, Wineberg S; The Renal Association. Clinical Practice Guideline: Undernutrition in Chronic Kidney Disease. Bristol (UK): The Renal Association; 2019. [Issued June 2019]. Available from: https://renal.org/wp-content/uploads/2019/06/FINAL-Nutrition-guideline-June-2019.pdf	Nutritional screening for improving professional practice for patient outcomes in hospital and primary care settings	10.1002/14651858.CD005539.pub2

<p>MacRobbie A, Johnston B, O'Neill C, Sives DA, Paterson E, Menzies K, O'Riordan M, Wilkinson M, Pryde N, Smithson R, Sweatn S, Boyce S, Johnston L; Healthcare Improvement Scotland. Scottish Palliative Care Guidelines. Edinburgh: Healthcare Improvement Scotland; 2019. Available from: https://www.palliativecareguidelines.scot.nhs.uk/</p>	Corticosteroids for the resolution of malignant bowel obstruction in advanced gynaecological and gastrointestinal cancer	10.1002/14651858.CD001219
	Surgery for the resolution of symptoms in malignant bowel obstruction in advanced gynaecological and gastrointestinal cancer	10.1002/14651858.CD002764
	Ketamine as an adjuvant to opioids for cancer pain	10.1002/14651858.CD003351
	Laxatives or methylnaltrexone for the management of constipation in palliative care patients	10.1002/14651858.CD003448.pub3
	Laxatives for the management of constipation in people receiving palliative care	10.1002/14651858.CD003448.pub4
	Methadone for cancer pain	10.1002/14651858.CD003971.pub3
	Exercise for the management of cancer-related fatigue in adults	10.1002/14651858.CD006145.pub3
	Medically assisted hydration for adult palliative care patients	10.1002/14651858.CD006273.pub3
	Interventions for the treatment of metastatic extradural spinal cord compression in adults	10.1002/14651858.CD006716.pub2
	Pharmacological treatments for fatigue associated with palliative care	10.1002/14651858.CD006788.pub2
	Droperidol for treatment of nausea and vomiting in palliative care patients	10.1002/14651858.CD006938.pub3
	Psychosocial interventions for reducing fatigue during cancer treatment in adults	10.1002/14651858.CD006953.pub2
	Benzodiazepines for the relief of breathlessness in advanced malignant and non-malignant diseases in adults	10.1002/14651858.CD007354.pub3
	Pharmacological interventions for pruritus in adult palliative care patients	10.1002/14651858.CD008320.pub3
	Topical lidocaine for neuropathic pain in adults	10.1002/14651858.CD010958.pub2
Opioids for cancer pain - an overview of Cochrane reviews	10.1002/14651858.CD012592.pub2	
<p>The British Pain Society. Position statement on the use of medical cannabis and cannabis-based products in the management of chronic pain. London: The British Pain Society; 2019. Available from: https://www.britishpainsociety.org/static/uploads/resources/files/BPS_position_statement_25_11_19.pdf</p>	Cannabis-based medicines for chronic neuropathic pain in adults	10.1002/14651858.CD012182.pub2

Trottier ED, Doré-Bergeron M-J, Chauvin-Kimoff L, Ali S; Acute Care Committee; Hospital Paediatrics Section; Community Paediatrics Section; Paediatric Emergency Medicine Section of the Canadian Paediatric Society. Managing pain and distress in children undergoing brief diagnostic and therapeutic procedures. Paediatr Child Health 2019 Nov;24(8):509-521. Available from: https://www.cps.ca/en/documents/position/managing-pain-and-distress	EMLA and Amethocaine for reduction of children's pain associated with needle insertion	10.1002/14651858.CD004236.pub2
	Psychological interventions for needle-related procedural pain and distress in children and adolescents	10.1002/14651858.CD005179.pub3
	Psychological interventions for needle-related procedural pain and distress in children and adolescents	10.1002/14651858.CD005179.pub4
	Non-pharmacological management of infant and young child procedural pain	10.1002/14651858.CD006275.pub3
	Sweet tasting solutions for reduction of needle-related procedural pain in children aged one to 16 years	10.1002/14651858.CD008408.pub3
	Sweet-tasting solutions for needle-related procedural pain in infants one month to one year of age	10.1002/14651858.CD008411.pub2
	Breastfeeding for procedural pain in infants beyond the neonatal period	10.1002/14651858.CD011248.pub2
Kulkarni GS, Black PC, Sridhar SS, Kapoor A, Zlotta AR, Shayegan B, Rendon RA, Chung P, van der Kwast T, Alimohamed N, Fradet Y, Kassouf W. Canadian Urological Association guideline: Muscle-invasive bladder cancer. Can Urol Assoc J. 2019 Aug;13(8):230-238. doi: 10.5489/cuaj.5902. Available from: https://www.ncbi.nlm.nih.gov/pmc/articles/PMC6737737/	Early palliative care for adults with advanced cancer	10.1002/14651858.CD011129.pub2
BC Centre for Palliative Care. B.C. Inter-professional Palliative Symptom Management Guidelines. New Westminster: BC Centre for Palliative Care; 2019. Available from: https://bc-cpc.ca/cpc/wp-content/uploads/2019/10/BCPCClinicalBestPractices-PrintVersionGreyscaleFull.pdf	Laxatives or methylnaltrexone for the management of constipation in palliative care patients	10.1002/14651858.CD003448.pub3
	Laxatives for the management of constipation in people receiving palliative care	10.1002/14651858.CD003448.pub4
	Oral morphine for cancer pain	10.1002/14651858.CD003868.pub4
	Oxycodone for cancer-related pain	10.1002/14651858.CD003870.pub5
	Megestrol acetate for treatment of anorexia-cachexia syndrome	10.1002/14651858.CD004310.pub3
	Eicosapentaenoic acid (EPA, an omega-3 fatty acid from fish oils) for the treatment of cancer cachexia	10.1002/14651858.CD004597.pub2
	Drug therapy for delirium in terminally ill adult patients	10.1002/14651858.CD004770.pub2
	Interventions for noisy breathing in patients near to death	10.1002/14651858.CD005177.pub2
	Non-pharmacological interventions for breathlessness in advanced stages of malignant and non-malignant diseases	10.1002/14651858.CD005623.pub2
Exercise for the management of cancer-related fatigue in adults	10.1002/14651858.CD006145.pub3	

	Codeine, alone and with paracetamol (acetaminophen), for cancer pain	10.1002/14651858.CD006601.pub4
	Pharmacological treatments for fatigue associated with palliative care	10.1002/14651858.CD006788.pub3
	Psychosocial interventions for reducing fatigue during cancer treatment in adults	10.1002/14651858.CD006953.pub2
	Benzodiazepines for the relief of breathlessness in advanced malignant and non-malignant diseases in adults	10.1002/14651858.CD007354.pub2
	Pharmacological interventions for pruritus in adult palliative care patients	10.1002/14651858.CD008320.pub2
	Interventions for fatigue and weight loss in adults with advanced progressive illness	10.1002/14651858.CD008427.pub2
	Interventions for treating persistent and intractable hiccups in adults	10.1002/14651858.CD008768.pub2
	Palliative pharmacological sedation for terminally ill adults	10.1002/14651858.CD010206.pub2
	Exercise for cancer cachexia in adults	10.1002/14651858.CD010804.pub2
	Opioids for the palliation of refractory breathlessness in adults with advanced disease and terminal illness	10.1002/14651858.CD011008.pub2
	Impact of morphine, fentanyl, oxycodone or codeine on patient consciousness, appetite and thirst when used to treat cancer pain	10.1002/14651858.CD011056.pub2
	Venlafaxine for neuropathic pain in adults	10.1002/14651858.CD011091.pub2
	Hydromorphone for cancer pain	10.1002/14651858.CD011108.pub2
	Psychosocial interventions for fatigue during cancer treatment with palliative intent	10.1002/14651858.CD012030
Alberta Provincial Gastrointestinal and Palliative Care Tumour Team, Alberta Health Services. Integrating an Early Palliative Approach into Advanced Colorectal Cancer Care (Clinical practice guideline supp-013) Version 1. Edmonton (Alberta): Alberta Health Services; 2019. [Issued January 2019]. Available from: https://www.albertahealthservices.ca/assets/info/hp/cancer/if-hp-cancer-guide-metastatic-colorectal-early-palliative-cancer-care.pdf	Early palliative care for adults with advanced cancer	10.1002/14651858.CD011129.pub2

AIM Specialty Health. Clinical Appropriateness Guidelines: Interventional Pain Management. Chicago, IL: AIM Specialty Health; 2019 January. Available from: http://www.aimspecialtyhealth.com/PDF/Guidelines/2019/Jan01/ AIM_Guidelines_MSK_Interventional-Pain-Management.pdf	Local anaesthetic sympathetic blockade for complex regional pain syndrome	10.1002/14651858.CD004598.pub4
American Diabetes Association. Standards of Medical Care in Diabetes - Position Statement 11: Microvascular Complications and Foot Care. Diabetes Care. 2019 Jan;42(Supplement 1):S124-S138. doi: 10.2337/dc19-S011. Available from: http://care.diabetesjournals.org/content/42/Supplement_1/S124	Pregabalin for acute and chronic pain in adults	10.1002/14651858.CD007076.pub2
	Gabapentin for chronic neuropathic pain in adults	10.1002/14651858.CD007938.pub4
Oskoui M, Pringsheim T, Billingham L, Potrebic S, Gersz EM, Gloss D, Holler-Managan Y, Leininger E, Licking N, Mack K, Powers SW, Sowell M, Victorio MC, Yonker M, Zanitsch H, Hershey AD. Practice guideline update summary: Pharmacologic treatment for pediatric migraine prevention: Report of the Guideline Development, Dissemination, and Implementation Subcommittee of the American Academy of Neurology and the American Headache Society. Neurology. 2019 Sep;93(11):500-509. doi: 10.1212/WNL.00000000000008105. Available from: https://www.ncbi.nlm.nih.gov/pubmed/31413170	Psychological therapies for the management of chronic and recurrent pain in children and adolescents	10.1002/14651858.CD003968.pub4
Hooten M, Thorson D, Hora J, Johnson C, Bianco J, Schweim K, Walker N, Clavel Jr A, Noonan MP, Reznikoff C, Bonte B, Wainio J, Kirksson E, Hansen A, Dvorkin J, Hadzik S. Health Care Guideline: Pain: Assessment, Non-Opioid Treatment Approaches and Opioid Management Care for Adults. Bloomington (MN): Institute for Clinical Systems Improvement (ICSI); 2019. [Issued September 2019]. Available from: https://www.icsi.org/wp-content/uploads/2019/10/Pain-Interactive-7th-V2-Ed-8.17.pdf	Lamotrigine for acute and chronic pain	10.1002/14651858.CD006044.pub2
	Long-term opioid management for chronic noncancer pain	10.1002/14651858.CD006605.pub2
	Phenytoin for neuropathic pain and fibromyalgia in adults	10.1002/14651858.CD009485.pub2
	Topical capsaicin (low concentration) for chronic neuropathic pain in adults	10.1002/14651858.CD010111
	Antiepileptic drugs for neuropathic pain and fibromyalgia - an overview of Cochrane reviews	10.1002/14651858.CD010567.pub2
	Oxycodone for neuropathic pain and fibromyalgia in adults	10.1002/14651858.CD010692.pub2
	Ketamine as an adjuvant to opioids for cancer pain	10.1002/14651858.CD003351.pub3
Swarm RA, Paice JA, Angheliescu DL, Are M, Bruce JY, Buga S, Chwistek M, Cleeland C, Craig D, Gafford E, Greenlee H, Kamal AH, Kamdar MM, LeGrand S, Mackey S, McDowell MR, Moryl N, Nabell LM, Nesbit S, Rabow MW, Rickerson E, Shatsky R, Sindt J, Urba SG, Youngwerth JM; National Comprehensive Cancer Network. NCCN	Oxycodone for cancer-related pain	10.1002/14651858.CD003870.pub6
	Methadone for cancer pain	10.1002/14651858.CD003971.pub4
	Opioids for the management of breakthrough pain in cancer patients	10.1002/14651858.CD004311.pub3

Clinical Practice Guidelines in Oncology: Adult Cancer Pain, Version 3.2019. Plymouth (PA): National Comprehensive Cancer Network; 2019 June. Available from: https://www.nccn.org/professionals/physician_gls/pdf/pain.pdf	Antidepressants for neuropathic pain	10.1002/14651858.CD005454
	Celiac plexus block for pancreatic cancer pain in adults	10.1002/14651858.CD007519.pub2
	Transdermal fentanyl for cancer pain	10.1002/14651858.CD010270.pub2
	Corticosteroids for the management of cancer-related pain in adults	10.1002/14651858.CD010756.pub2
	Hydromorphone for cancer pain	10.1002/14651858.CD011108.pub2
	Tramadol with or without paracetamol (acetaminophen) for cancer pain	10.1002/14651858.CD012508.pub2
Denlinger CS, Sanft T, Armenian S, Baker KS, Broderick G, Demark-Wahnefried W, Friedman DL, Goldman M, Hudson M, Khakpour N, Koura D, Lally RM, Langbaum TS, McDonough AL, Melisko M, Mooney K, Moore HCF, Moslehi JJ, O'Connor T, Overholser L, Paskett ED, Peterson L, Pirl W, Rodriguez A, Ruddy KJ, Smith S, Syrjala KL, Tevaarwerk A, Urba SG, Zee P; National Comprehensive Cancer Network. NCCN Clinical Practice Guidelines in Oncology: Survivorship, Version 2.2019. Plymouth (PA): National Comprehensive Cancer Network; 2019 June. Available from: https://www.nccn.org/professionals/physician_gls/pdf/survivorshi	Exercise for the management of cancer-related fatigue in adults	10.1002/14651858.CD006145.pub3
	Transcutaneous electric nerve stimulation (TENS) for cancer pain in adults	10.1002/14651858.CD006276.pub3
	Pharmacological treatments for fatigue associated with palliative care	10.1002/14651858.CD006788.pub3
	Gabapentin for chronic neuropathic pain and fibromyalgia in adults	10.1002/14651858.CD007938.pub2
Clinical Oncology Society of Australia. COSA Position Statement on Exercise in Cancer Care. Sydney: Clinical Oncology Society of Australia; 2018. [Version 1a Issued October 2019]. Available from: https://www.cosa.org.au/media/332583/cosa-position-statement-oct2019-web-final.pdf	Exercise for the management of cancer-related fatigue in adults	10.1002/14651858.CD006145.pub3
Yang IA, Dabscheck E, George J, Jenkins S, McDonald CF, McDonald V, Smith B, Zwar N; on behalf of the Lung Foundation Australia and the Thoracic Society of Australia and New Zealand. The COPD-X Plan: Australian and New Zealand Guidelines for the management of Chronic Obstructive Pulmonary Disease 2019. Version 2.59, August 2019. Sydney, NSW, Australia: Lung Foundation Australia and Thoracic Society of Australia and New Zealand; 2019. Available from: https://copdx.org.au/wp-content/uploads/2019/11/COPDX-V2-59-Aug-2019-FINAL2.pdf	Benzodiazepines for the relief of breathlessness in advanced malignant and non-malignant diseases in adults	10.1002/14651858.CD007354.pub3
	Opioids for the palliation of refractory breathlessness in adults with advanced disease and terminal illness	10.1002/14651858.CD011008.pub2
Global Initiative for Chronic Obstructive Lung Disease. Global strategy for the diagnosis, management and prevention of chronic	Non-pharmacological interventions for breathlessness in advanced stages of malignant and non-malignant diseases	10.1002/14651858.CD005623.pub2

obstructive pulmonary disease: 2019 report. Vancouver (WA): Global Initiative for Chronic Obstructive Lung Disease (GOLD); 2019. Available from: https://goldcopd.org/wp-content/uploads/2018/11/GOLD-2019-v1.7-FINAL-14Nov2018-WMS.pdf	Benzodiazepines for the relief of breathlessness in advanced malignant and non-malignant diseases in adults	10.1002/14651858.CD007354.pub2
	Interventions for fatigue and weight loss in adults with advanced progressive illness	10.1002/14651858.CD008427.pub2
Dansk Selskab for Anaesthesiologi og Intensiv Medicin. National klinisk retningslinje: Systematisk opfølgning af patienter i behandling med opioider for non-maligne smerter. [National Clinical Guideline on systematic follow-up of patients treated with opioids for non-cancer pain]. Copenhagen: Sundhedsstyrelsen, Danish Health Authority; June 2019. Available from: https://www.sst.dk/-/media/Udgivelser/2019/published_guideline_3348-1_0.ashx?la=da&hash=6F5B2089A38DBF1ADD75088C73CF984277ABA7B5	Adverse events associated with medium- and long-term use of opioids for chronic non-cancer pain: an overview of Cochrane Reviews	10.1002/14651858.CD012509.pub2
Gesellschaft für Neuropädiatrie (GNP). S3 Leitlinie: Diagnose und Therapie des Guillain-Barré Syndroms im Kindes- und Jugendalter (ICD-10: G61.0). [S3 Guideline: Diagnosis and Treatment of Guillain-Barré Syndrome in Childhood and Adolescence (ICD-10: G61.0)]. Kempen: Gesellschaft für Neuropädiatrie (GNP); 2019 March. Available from: https://www.awmf.org/uploads/tx_szleitlinien/022-008l_S3_Guillain-Barre_Syndrom_2019-03.pdf	Gabapentin for acute and chronic pain	10.1002/14651858.CD005452
Bendtsen L, Zakrzewska JM, Abbott J, Braschinsky M, Di Stefano G, Donnet A, Eide PK, Leal PRL, Maarbjerg S, May A, Nurmikko T, Obermann M, Jensen TS, Cruccu G. European Academy of Neurology guideline on trigeminal neuralgia. Eur J Neurol. 2019 Jun;26(6):831-849. doi: 10.1111/ene.13950. Available from: https://onlinelibrary.wiley.com/doi/full/10.1111/ene.13950	Carbamazepine for chronic neuropathic pain and fibromyalgia in adults	10.1002/14651858.CD005451.pub3
	Psychological therapies for the management of chronic pain (excluding headache) in adults	10.1002/14651858.CD007407.pub3
Deutsche Gesellschaft für Neurologie (DGN). S2k Leitlinien für Diagnostik und Therapie in der Neurologie: Diagnose und nicht interventionelle Therapie neuropathischer Schmerzen. [S2k Guidelines for diagnosis and treatment in neurology: diagnosis and non-interventional treatment of neuropathic pain]. Berlin:	Carbamazepine for chronic neuropathic pain and fibromyalgia in adults	10.1002/14651858.CD005451.pub3
	Opioids for neuropathic pain	10.1002/14651858.CD006146.pub2
	Topical capsaicin (high concentration) for chronic neuropathic pain in adults	10.1002/14651858.CD007393.pub4

Deutsche Gesellschaft für Neurologie (DGN); 2019. Available from: https://www.awmf.org/uploads/tx_szleitlinien/030-114l_S2k_Diagnose-nicht-interventionelle-Therapie-neuropathischer-Schmerzen_2019-09.pdf

Gabapentin for chronic neuropathic pain and fibromyalgia in adults	10.1002/14651858.CD007938.pub3
Gabapentin for chronic neuropathic pain in adults	10.1002/14651858.CD007938.pub4
Amitriptyline for neuropathic pain in adults	10.1002/14651858.CD008242.pub3
Combination pharmacotherapy for the treatment of neuropathic pain in adults	10.1002/14651858.CD008943.pub2
Phenytoin for neuropathic pain and fibromyalgia in adults	10.1002/14651858.CD009485.pub2
Clonazepam for neuropathic pain and fibromyalgia in adults	10.1002/14651858.CD009486.pub2
Antiepileptic drugs for neuropathic pain and fibromyalgia - an overview of Cochrane reviews	10.1002/14651858.CD010567.pub2
Oxycodone for neuropathic pain in adults	10.1002/14651858.CD010692.pub3
Imipramine for neuropathic pain in adults	10.1002/14651858.CD010769.pub2
Oral nonsteroidal anti-inflammatory drugs for neuropathic pain	10.1002/14651858.CD010902.pub2
Topical lidocaine for neuropathic pain in adults	10.1002/14651858.CD010958.pub2
Venlafaxine for neuropathic pain in adults	10.1002/14651858.CD011091.pub2
Nortriptyline for neuropathic pain in adults	10.1002/14651858.CD011209.pub2
Psychological therapies for the management of chronic neuropathic pain in adults	10.1002/14651858.CD011259.pub2
Buprenorphine for neuropathic pain in adults	10.1002/14651858.CD011603.pub2
Hydromorphone for neuropathic pain in adults	10.1002/14651858.CD011604.pub2
Fentanyl for neuropathic pain in adults	10.1002/14651858.CD011605.pub2
Morphine for chronic neuropathic pain in adults	10.1002/14651858.CD011669.pub2
Cannabis-based medicines for chronic neuropathic pain in adults	10.1002/14651858.CD012182.pub2
Paracetamol (acetaminophen) with or without codeine or dihydrocodeine for neuropathic pain in adults	10.1002/14651858.CD012227.pub2
Methadone for neuropathic pain in adults	10.1002/14651858.CD012499.pub2
Milnacipran for neuropathic pain in adults	10.1002/14651858.CD011789
Transcutaneous Electrical Nerve Stimulation (TENS) for neuropathic pain in adults	10.1002/14651858.CD011976
Eicosapentaenoic acid (EPA, an omega-3 fatty acid from fish oils) for the treatment of cancer cachexia	10.1002/14651858.CD004597.pub2
Homeopathic medicines for adverse effects of cancer treatments	10.1002/14651858.CD004845.pub2

Leitlinienprogramm Onkologie (Deutsche Krebsgesellschaft; Deutsche Krebshilfe; AWMF). S3 Leitlinie Magenkarzinom: Diagnostik und Therapie der Adenokarzinome des Magens und ösophagogastralen Übergangs. [S3 Guideline Gastric carcinoma:

diagnosis and treatment of gastric adenocarcinoma and esophagogastral junction]. Berlin: Leitlinienprogramm Onkologie; 2019. Available from: https://www.awmf.org/leitlinien/detail/II/032-009OL.html	Selenium for alleviating the side effects of chemotherapy, radiotherapy and surgery in cancer patients	10.1002/14651858.CD005037.pub2
	Psychosocial interventions for reducing fatigue during cancer treatment in adults	10.1002/14651858.CD006953.pub2
	Acupuncture for cancer pain in adults	10.1002/14651858.CD007753.pub2
	Massage with or without aromatherapy for symptom relief in people with cancer	10.1002/14651858.CD009873.pub3
	Exercise for cancer cachexia in adults	10.1002/14651858.CD010804.pub2
Leitlinienprogramm Onkologie (Deutsche Krebsgesellschaft; Deutsche Krebshilfe; AWMF). S3-Leitlinie: Palliativmedizin für Patienten mit einer nicht heilbaren krebserkrankung. [S3 guideline: palliative care for patients with incurable cancer]. Berlin: Leitlinienprogramm Onkologie; 2019. Available from: https://www.leitlinienprogramm-onkologie.de/fileadmin/user_upload/Downloads/Leitlinien/Palliativmedizin/Version_2/LL_Palliativmedizin_2.0_Langversion.pdf	Corticosteroids for the resolution of malignant bowel obstruction in advanced gynaecological and gastrointestinal cancer	10.1002/14651858.CD001219
	Opioids for the palliation of breathlessness in terminal illness	10.1002/14651858.CD002066
	Aromatherapy and massage for symptom relief in patients with cancer	10.1002/14651858.CD002287.pub2
	Surgery for the resolution of symptoms in malignant bowel obstruction in advanced gynaecological and gastrointestinal cancer	10.1002/14651858.CD002764.pub2
	Benzodiazepines and related drugs for insomnia in palliative care	10.1002/14651858.CD003346
	Hydromorphone for acute and chronic pain	10.1002/14651858.CD003447
	Laxatives or methylnaltrexone for the management of constipation in palliative care patients	10.1002/14651858.CD003448.pub3
	Laxatives for the management of constipation in people receiving palliative care	10.1002/14651858.CD003448.pub4
	Oral morphine for cancer pain	10.1002/14651858.CD003868.pub2
	Oral morphine for cancer pain	10.1002/14651858.CD003868.pub3
	Methadone for cancer pain	10.1002/14651858.CD003971.pub3
	Megestrol acetate for treatment of anorexia-cachexia syndrome	10.1002/14651858.CD004310.pub3
	Opioids for the management of breakthrough (episodic) pain in cancer patients	10.1002/14651858.CD004311.pub2
	Opioids for the management of breakthrough pain in cancer patients	10.1002/14651858.CD004311.pub3
	Drug therapy for symptoms associated with anxiety in adult palliative care patients	10.1002/14651858.CD004596.pub3

Oxygen therapy for dyspnoea in adults	10.1002/14651858.CD004769.pub2
Drug therapy for delirium in terminally ill adult patients	10.1002/14651858.CD004770.pub2
Opioid switching to improve pain relief and drug tolerability	10.1002/14651858.CD004847
Interventions for noisy breathing in patients near to death	10.1002/14651858.CD005177.pub2
NSAIDs or paracetamol, alone or combined with opioids, for cancer pain	10.1002/14651858.CD005180
Psychotherapy for depression among incurable cancer patients	10.1002/14651858.CD005537.pub2
Non-pharmacological interventions for breathlessness in advanced stages of malignant and non-malignant diseases	10.1002/14651858.CD005623.pub2
Haloperidol for the treatment of nausea and vomiting in palliative care patients	10.1002/14651858.CD006271.pub3
Medically assisted hydration for adult palliative care patients	10.1002/14651858.CD006273.pub3
Mu-opioid antagonists for opioid-induced bowel dysfunction in people with cancer and people receiving palliative care	10.1002/14651858.CD006332.pub3
Pharmacological treatments for fatigue associated with palliative care	10.1002/14651858.CD006788.pub3
Droperidol for treatment of nausea and vomiting in palliative care patients	10.1002/14651858.CD006938.pub3
Psychosocial interventions for reducing fatigue during cancer treatment in adults	10.1002/14651858.CD006953.pub2
Benzodiazepines for the relief of breathlessness in advanced malignant and non-malignant diseases in adults	10.1002/14651858.CD007354.pub3
Effectiveness and cost-effectiveness of home palliative care services for adults with advanced illness and their caregivers	10.1002/14651858.CD007760.pub2
Educational interventions for the management of cancer-related fatigue in adults	10.1002/14651858.CD008144.pub2
Interventions for fatigue and weight loss in adults with advanced progressive illness	10.1002/14651858.CD008427.pub2
Levomepromazine for nausea and vomiting in palliative care	10.1002/14651858.CD009420.pub3
Transdermal fentanyl for cancer pain	10.1002/14651858.CD010270.pub2

	Opioids for the palliation of refractory breathlessness in adults with advanced disease and terminal illness	10.1002/14651858.CD011008.pub2
	Early palliative care for adults with advanced cancer	10.1002/14651858.CD011129.pub2
	Corticosteroids for adult patients with advanced cancer who have nausea and vomiting (not related to chemotherapy, radiotherapy, or surgery)	10.1002/14651858.CD012002.pub2
	Psychosocial interventions for fatigue during cancer treatment with palliative intent	10.1002/14651858.CD012030.pub2
	Opioids for cancer pain - an overview of Cochrane reviews	10.1002/14651858.CD012592.pub2
	Systemic corticosteroids for the management of cancer-related breathlessness (dyspnoea) in adults	10.1002/14651858.CD012704.pub2
Suomalaisen Lääkäriseuran Duodecimin, Suomen Palliatiivisen Lääketieteen yhdistyksen asettama työryhmä, Finnish Medical Society Duodecim, Finnish Palliative Medicine Association. Duodecim Käypä Hoito: Palliatiivinen hoito ja saattohoito. [Duodecim Current Care Guidelines: Palliative care and end-of-life care]. Helsinki: Finnish Medical Society Duodecim; October 2019. Available from: https://www.kaypahoito.fi/hoi50063	Corticosteroids for the resolution of malignant bowel obstruction in advanced gynaecological and gastrointestinal cancer	10.1002/14651858.CD001219
	Opioids for the palliation of breathlessness in terminal illness	10.1002/14651858.CD002066
	Bisphosphonates for the relief of pain secondary to bone metastases	10.1002/14651858.CD002068
	Pleurodesis for malignant pleural effusions	10.1002/14651858.CD002916.pub2
	Calcitonin for metastatic bone pain	10.1002/14651858.CD003223.pub2
	Radioisotopes for metastatic bone pain	10.1002/14651858.CD003347.pub2
	Ketamine as an adjuvant to opioids for cancer pain	10.1002/14651858.CD003351.pub2
	Laxatives or methylnaltrexone for the management of constipation in palliative care patients	10.1002/14651858.CD003448.pub3
	Laxatives for the management of constipation in people receiving palliative care	10.1002/14651858.CD003448.pub4
	Oral morphine for cancer pain	10.1002/14651858.CD003868.pub4
	Oxycodone for cancer-related pain	10.1002/14651858.CD003870.pub5
	Methadone for cancer pain	10.1002/14651858.CD003971.pub3
	Megestrol acetate for the treatment of anorexia-cachexia syndrome	10.1002/14651858.CD004310.pub2
	Megestrol acetate for treatment of anorexia-cachexia syndrome	10.1002/14651858.CD004310.pub3
	Opioids for the management of breakthrough (episodic) pain in cancer patients	10.1002/14651858.CD004311.pub2

Opioids for the management of breakthrough pain in cancer patients	10.1002/14651858.CD004311.pub3
Drug therapy for symptoms associated with anxiety in adult palliative care patients	10.1002/14651858.CD004596.pub2
Oxygen therapy for dyspnoea in adults	10.1002/14651858.CD004769.pub2
Drug therapy for delirium in terminally ill adult patients	10.1002/14651858.CD004770.pub2
Interventions for noisy breathing in patients near to death	10.1002/14651858.CD005177.pub2
Antidepressants for neuropathic pain	10.1002/14651858.CD005454.pub2
Psychotherapy for depression among incurable cancer patients	10.1002/14651858.CD005537.pub2
Non-pharmacological interventions for breathlessness in advanced stages of malignant and non-malignant diseases	10.1002/14651858.CD005623.pub2
Haloperidol for the treatment of nausea and vomiting in palliative care patients	10.1002/14651858.CD006271.pub2
Medically assisted hydration for adult palliative care patients	10.1002/14651858.CD006273.pub3
Medically assisted nutrition for palliative care in adult patients	10.1002/14651858.CD006274.pub2
Pharmacologic interventions for treating phantom limb pain	10.1002/14651858.CD006380.pub3
Codeine, alone and with paracetamol (acetaminophen), for cancer pain	10.1002/14651858.CD006601.pub4
Pharmacological treatments for fatigue associated with palliative care	10.1002/14651858.CD006788.pub3
Benzodiazepines for the relief of breathlessness in advanced malignant and non-malignant diseases in adults	10.1002/14651858.CD007354.pub3
Topical capsaicin (high concentration) for chronic neuropathic pain in adults	10.1002/14651858.CD007393.pub4
Gabapentin for chronic neuropathic pain in adults	10.1002/14651858.CD007938.pub4
Combination pharmacotherapy for the treatment of neuropathic pain in adults	10.1002/14651858.CD008943.pub2
Blood transfusions for anaemia in patients with advanced cancer	10.1002/14651858.CD009007.pub2
Spinal cord stimulation for cancer-related pain in adults	10.1002/14651858.CD009389.pub3
Palliative pharmacological sedation for terminally ill adults	10.1002/14651858.CD010206.pub2

	Antiepileptic drugs for neuropathic pain and fibromyalgia - an overview of Cochrane reviews	10.1002/14651858.CD010567.pub2
	Opioids for the palliation of refractory breathlessness in adults with advanced disease and terminal illness	10.1002/14651858.CD011008.pub2
	Venlafaxine for neuropathic pain in adults	10.1002/14651858.CD011091.pub2
	Tramadol with or without paracetamol (acetaminophen) for cancer pain	10.1002/14651858.CD012508.pub2
	Oral paracetamol (acetaminophen) for cancer pain	10.1002/14651858.CD012637.pub2
	Oral nonsteroidal anti-inflammatory drugs (NSAIDs) for cancer pain in adults	10.1002/14651858.CD012638.pub2
Suomalaisen Lääkäriseuran Duodecimin, Suomen Keuhkolääkäriyhdistys ry:n asettama työryhmä, Finnish Medical Society Duodecim, Finnish Association of Respiratory Physicians. Duodecim Käypä Hoito: Keuhkohtaumatauti. [Duodecim Current Care Guidelines: Chronic Obstructive Pulmonary Disease]. Helsinki: Finnish Medical Society Duodecim; November 2019. Available from: https://www.kaypahoito.fi/hoi06040	Opioids for the palliation of breathlessness in terminal illness	10.1002/14651858.CD002066
	Opioids for the palliation of refractory breathlessness in adults with advanced disease and terminal illness	10.1002/14651858.CD011008.pub2
Colombo N, Sessa C, du Bois A, Ledermann J, McCluggage WG, McNeish I, Morice P, Pignata S, Ray-Coquard I, Vergote I, Baert T, Belaroussi I, Dashora A, Olbrecht S, Planchamp F, Querleu D; ESMO-ESGO Ovarian Cancer Consensus Conference Working Group. ESMO-ESGO consensus conference recommendations on ovarian cancer: pathology and molecular biology, early and advanced stages, borderline tumours and recurrent disease. Ann Oncol 2019 May;30(5):672-705. doi: 10.1093/annonc/mdz062. Available from: https://www.ncbi.nlm.nih.gov/pubmed/31046081	Corticosteroids for the resolution of malignant bowel obstruction in advanced gynaecological and gastrointestinal cancer	10.1002/14651858.CD001219
	Surgery for the resolution of symptoms in malignant bowel obstruction in advanced gynaecological and gastrointestinal cancer	10.1002/14651858.CD002764.pub2
	Early palliative care for adults with advanced cancer	10.1002/14651858.CD011129.pub2
Engeler D, Baranowski AP, Borovicka J, Berghmans B, Cottrell A, Elneil S, Hughes J, Messelink EJ, de C Williams AC; European Association of Urology . Guidelines on Chronic Pelvic Pain. Arnhem, The Netherlands: European Association of Urology; 2019. [Updated 2019]. Available from: https://uroweb.org/guideline/chronic-pelvic-pain/	Transcutaneous electrical nerve stimulation (TENS) for chronic pain	10.1002/14651858.CD003222.pub2
	Carbamazepine for acute and chronic pain in adults	10.1002/14651858.CD005451.pub2
	Antidepressants for neuropathic pain	10.1002/14651858.CD005454.pub2
	Long-term opioid management for chronic noncancer pain	10.1002/14651858.CD006605.pub2
	Pregabalin for acute and chronic pain in adults	10.1002/14651858.CD007076.pub2

	Psychological therapies for the management of chronic pain (excluding headache) in adults	10.1002/14651858.CD007407.pub3
	Gabapentin for chronic neuropathic pain and fibromyalgia in adults	10.1002/14651858.CD007938.pub2
Kouremenos E, Arvaniti C, Constantinidis TS, Giannouli E, Fakas N, Kalamatas T, Kararizou E, Naoumis D, Mitsikostas DD; Hellenic Headache Society. Consensus of the Hellenic Headache Society on the diagnosis and treatment of migraine. J Headache Pain 2019 Dec;20(1):113. doi: 10.1186/s10194-019-1060-6. Available from: https://www.ncbi.nlm.nih.gov/pubmed/31835997	Aspirin with or without an antiemetic for acute migraine headaches in adults	10.1002/14651858.CD008041.pub3
Wainwright TW, Gill M, McDonald DA, Middleton RG, Reed M, Sahota O, Yates P, Ljungqvist O. Consensus statement for perioperative care in total hip replacement and total knee replacement surgery: Enhanced Recovery After Surgery (ERAS®) Society recommendations. Acta Orthop 2019 Oct. doi: 10.1080/17453674.2019.1683790. [Epub ahead of print]. Available from: https://www.ncbi.nlm.nih.gov/pubmed/31663402	Epidural analgesia for pain relief following hip or knee replacement	10.1002/14651858.CD003071
Gustafsson UO, Scott MJ, Hubner M, Nygren J, Demartines N, Francis N, Rockall TA, Young-Fadok TM, Hill AG, Soop M, de Boer HD, Urman RD, Chang GJ, Fichera A, Kessler H, Grass F, Whang EE, Fawcett WJ, Carli F, Lobo DN, Rollins KE, Balfour A, Baldini G, Riedel B, Ljungqvist O. Guidelines for Perioperative Care in Elective Colorectal Surgery: Enhanced Recovery After Surgery (ERAS®) Society Recommendations: 2018. World J Surg 2019 Mar;43(3):659-695. doi: 10.1007/s00268-018-4844-y. Available from: https://www.ncbi.nlm.nih.gov/pubmed/30426	Perioperative ketamine for acute postoperative pain	10.1002/14651858.CD004603.pub2
	Perioperative transversus abdominis plane (TAP) blocks for analgesia after abdominal surgery	10.1002/14651858.CD007705.pub2
	Psychological interventions for acute pain after open heart surgery	10.1002/14651858.CD009984.pub3
	Liposomal bupivacaine infiltration at the surgical site for the management of postoperative pain	10.1002/14651858.CD011419.pub2

<p>Low DE, Allum W, De Manzoni G, Ferri L, Immanuel A, Kuppusamy M, Law S, Lindblad M, Maynard N, Neal J, Pramesh CS, Scott M, Mark Smithers B, Addor V, Ljungqvist O. Guidelines for Perioperative Care in Esophagectomy: Enhanced Recovery After Surgery (ERAS®) Society Recommendations. World J Surg 2019 Feb;43(2):299-330. doi: 10.1007/s00268-018-4786-4. Available from: https://www.ncbi.nlm.nih.gov/pubmed/30276441</p>	<p>Single dose oral paracetamol (acetaminophen) for postoperative pain</p>	<p>10.1002/14651858.CD004602.pub2</p>
<p>Schönhofer B, Geiseler J, Braune S, Dellweg D, Fuchs H, Hirschfeld-Araujo J, Janssens U, Mörer O, Rollnik J, Rosseau S, Schreiter D, Weber-Carstens S, Windisch W, Westhoff M; Deutsche Gesellschaft für Pneumologie und Beatmungsmedizin e.V. (DGP). S2k-Leitlinie: Prolonged Weaning. [S2k-guideline: Prolonged Weaning]. Berlin: Deutsche Gesellschaft für Pneumologie und Beatmungsmedizin e.V. (DGP); 2019 August. Available from: https://www.awmf.org/uploads/tx_szleitlinien/020-015l_S2k_Prolongiertes_Weaning_2019_09.pdf</p>	<p>Neuromuscular electrical stimulation for muscle weakness in adults with advanced disease</p>	<p>10.1002/14651858.CD009419.pub3</p>
<p>Gross GE; Deutsche Dermatologische Gesellschaft (DDG); Paul-Ehrlich-Gesellschaft für Chemotherapie e.V. (PEG). S2k-Leitlinie: Diagnostik und Therapie des Zoster und der Postzosterneuralgie. [S2k-guideline: Diagnosis and treatment of herpetic and post-herpetic neuralgia]. Rheinbach: Paul-Ehrlich-Gesellschaft für Chemotherapie e.V. (PEG); 2019 May. Available from: https://www.awmf.org/leitlinien/detail/II/013-023.html.</p>	<p>Topical lidocaine for the treatment of postherpetic neuralgia</p>	<p>10.1002/14651858.CD004846.pub2</p>
	<p>Topical capsaicin (high concentration) for chronic neuropathic pain in adults</p>	<p>10.1002/14651858.CD007393.pub4</p>
	<p>Gabapentin for chronic neuropathic pain in adults</p>	<p>10.1002/14651858.CD007938.pub4</p>
	<p>Topical lidocaine for neuropathic pain in adults</p>	<p>10.1002/14651858.CD010958.pub2</p>
<p>Evcik D, Ketenci A, Sindel D. The Turkish Society of Physical Medicine and Rehabilitation (TSPMR) guideline recommendations for the management of fibromyalgia syndrome. Turk J Phys Med Rehabil. 2019 May;65(2):111-123. doi: 10.5606/tftrd.2019.4815. Available from: https://www.ncbi.nlm.nih.gov/pubmed/31453551</p>	<p>Amitriptyline for neuropathic pain and fibromyalgia in adults</p>	<p>10.1002/14651858.CD008242.pub2</p>
	<p>Physical activity and exercise for chronic pain in adults: an overview of Cochrane Reviews</p>	<p>10.1002/14651858.CD011279.pub3</p>
	<p>Gabapentin for fibromyalgia pain in adults</p>	<p>10.1002/14651858.CD012188.pub2</p>
	<p>Oral nonsteroidal anti-inflammatory drugs for fibromyalgia in adults</p>	<p>10.1002/14651858.CD012332.pub2</p>

<p>Bundesärztekammer; Kassenärztliche Bundesvereinigung; Arbeitsgemeinschaft der Wissenschaftlichen Medizinischen Fachgesellschaften. Nationale VersorgungsLeitlinie: Chronische Herzinsuffizienz (3. Auflage). [National Clinical Guideline: Chronic heart failure (3rd edition)]. Berlin: Ärztliches Zentrum für Qualität in der Medizin; 2019. Available from: https://www.awmf.org/uploads/tx_szleitlinien/nvl-006l_S3_Chronische_Herzinsuffizienz_2019-10_01.pdf</p>	<p>Effectiveness and cost-effectiveness of home palliative care services for adults with advanced illness and their caregivers</p>	<p>10.1002/14651858.CD007760.pub2</p>
<p>Filetti S, Durante C, Hartl D, Leboulleux S, Locati LD, Newbold K, Papotti MG, Berruti A; European Society for Medical Oncology (ESMO) Guidelines Committee. Thyroid cancer: ESMO Clinical Practice Guidelines for diagnosis, treatment and follow-up. <i>Annals of Oncology</i>. 2019 Sept: mdz400. doi: 10.1093/annonc/mdz400. Available from: https://www.esmo.org/Guidelines/Endocrine-and-Neuroendocrine-Cancers/Thyroid-cancer.</p>	<p>Interventions for the treatment of metastatic extradural spinal cord compression in adults</p>	<p>10.1002/14651858.CD006716.pub3</p>
<p>Deutsche Gesellschaft für Gynäkologie und Geburtshilfe (Deutsche Gesellschaft für Gynäkologie und Geburtshilfe (DGGG)). S3-Leitlinie Diagnostik: Therapie und Nachsorge maligner Ovarialtumoren. [S3 guideline diagnostics: Therapy and follow-up malignant ovarian tumors]. Berlin: Leitlinienprogramm Onkologie; 2019 January. Available from: https://www.leitlinienprogramm-onkologie.de/fileadmin/user_upload/Downloads/Leitlinien/Ovarialkarzinom/Version_3__2018_/LL_Ovarialkarzinom_Langversion_3.0.pdf</p>	<p>Pleurodesis for malignant pleural effusions Exercise for the management of cancer-related fatigue in adults</p>	<p>10.1002/14651858.CD002916.pub2 10.1002/14651858.CD006145.pub3</p>

<p>Oommen PT, Schütz C, Baltruschat K, Deuter C, Foeldvari I, Ganser G, Haas JP, Hinze C, Holzinger D, Hospach A, Huppertz HI, Illhardt A, Jung M, Kallinich T, Klein A, Minden K, Mönkemöller K, Mrusek S, Niehues T, Neudorf U, Schneider M, Schoof P, Thon A, Wachowsky M, Wagner N; Deutsche Gesellschaft für Kinder- und Jugendmedizin e.V. (DGKJ); Gesellschaft für Kinder- und Jugendrheumatologie e.V. (GKJR). S2k Leitlinie: Therapie der Juvenilen Idiopathischen Arthritis. [S2k Guideline: Treatment for Juvenile Idiopathic Arthritis]. Berlin: Deutsche Gesellschaft für Kinder- und Jugendmedizin e.V. (DGKJ); 2019 November. Available from: https://www.awmf.org/leitlinien/detail/II/027-020.html.</p>	<p>Non-steroidal anti-inflammatory drugs (NSAIDs) for chronic non-cancer pain in children and adolescents</p>	<p>10.1002/14651858.CD012537.pub2</p>
<p>Schröder HM, Lilienthal S, Schreiber-Gollwitzer BM, Grießmeier B, Hesselbarth B, Lein-Köhler I, Nest A, Weiler-Wichtl LJ, Leiss U; Gesellschaft für Pädiatrische Onkologie und Hämatologie (GPOH). S3-Leitlinie: Psychosoziale Versorgung in der Pädiatrischen Onkologie und Hämatologie. [S3-Guideline: Psychosocial care in paediatric oncology and haematology]. Berlin: Gesellschaft für Pädiatrische Onkologie und Hämatologie (GPOH); 2019. Available from: https://www.awmf.org/leitlinien/detail/II/025-002.html</p>	<p>Psychological therapies for the management of chronic and recurrent pain in children and adolescents</p>	<p>10.1002/14651858.CD003968</p>
<p>Ministry of Health Malaysia; Academy of Medicine Malaysia; Malaysian Psychiatric Association. Clinical Practice Guidelines: Management of Major Depressive Disorder (Second Edition). Putrajaya, Malaysia: Ministry of Health Malaysia; 2019. Available from: http://www.acadmed.org.my/index.cfm?&menuid=67</p>	<p>Amitriptyline for neuropathic pain in adults</p>	<p>10.1002/14651858.CD008242.pub3</p>
<p>Scottish Intercollegiate Guidelines Network (SIGN). Management of epithelial ovarian cancer. Edinburgh: SIGN; 2018. (SIGN publication no. 135). [Issued November 2013; last updated October 2018]. Available from: https://www.sign.ac.uk/assets/sign135_oct2018.pdf</p>	<p>Corticosteroids for the resolution of malignant bowel obstruction in advanced gynaecological and gastrointestinal cancer</p>	<p>10.1002/14651858.CD001219</p>
	<p>Surgery for the resolution of symptoms in malignant bowel obstruction in advanced gynaecological and gastrointestinal cancer</p>	<p>10.1002/14651858.CD002764</p>
<p>Scottish Intercollegiate Guidelines Network (SIGN). Pharmacological management of migraine. Edinburgh: SIGN; 2018.</p>	<p>Ibuprofen with or without an antiemetic for acute migraine headaches in adults</p>	<p>10.1002/14651858.CD008039.pub3</p>

(SIGN publication no. 155). [Issued February 2018]. Available from: <http://www.sign.ac.uk/assets/sign155.pdf>

Paracetamol (acetaminophen) with or without an antiemetic for acute migraine headaches in adults	10.1002/14651858.CD008040.pub3
Aspirin with or without an antiemetic for acute migraine headaches in adults	10.1002/14651858.CD008041.pub3
Sumatriptan plus naproxen for the treatment of acute migraine attacks in adults	10.1002/14651858.CD008541.pub3
Zolmitriptan for acute migraine attacks in adults	10.1002/14651858.CD008616.pub2
Diclofenac with or without an antiemetic for acute migraine headaches in adults	10.1002/14651858.CD008783.pub3
Sumatriptan (all routes of administration) for acute migraine attacks in adults - overview of Cochrane reviews	10.1002/14651858.CD009108.pub2
Naproxen with or without an antiemetic for acute migraine headaches in adults	10.1002/14651858.CD009455.pub2
Antiepileptics other than gabapentin, pregabalin, topiramate, and valproate for the prophylaxis of episodic migraine in adults	10.1002/14651858.CD010608
Selective serotonin reuptake inhibitors (SSRIs) and serotonin-norepinephrine reuptake inhibitors (SNRIs) for the prevention of tension-type headache in adults	10.1002/14651858.CD011681
Gabapentin or pregabalin for the prophylaxis of episodic migraine in adults	10.1002/14651858.CD010609
Topiramate for the prophylaxis of episodic migraine in adults	10.1002/14651858.CD010610
Valproate (valproic acid or sodium valproate or a combination of the two) for the prophylaxis of episodic migraine in adults	10.1002/14651858.CD010611
National Clinical Guideline Centre; National Institute for Health and Care Excellence (commissioner). Chronic obstructive pulmonary disease in over 16s: diagnosis and management. London: National Clinical Guideline Centre, Royal College of Physicians; 2018. (NICE NG115). [Issued December 2018]. Available from: https://www.nice.org.uk/guidance/ng115 .	Opioids for the palliation of breathlessness in terminal illness 10.1002/14651858.CD002066
National Guideline Centre (hosted by the Royal College of Physicians); National Institute for Health and Care Excellence (commissioner). Emergency and acute medical care in over 16s:	Effectiveness and cost-effectiveness of home palliative care services for adults with advanced illness and their caregivers 10.1002/14651858.CD007760.pub2

service delivery and organisation. London: National Institute for Health and Care Excellence; 2018. (NICE NG94). [Issued March 2018]. Available from: https://www.nice.org.uk/guidance/ng94	End-of-life care pathways for improving outcomes in caring for the dying	10.1002/14651858.CD008006.pub3
	Advance care planning for end-of-life care	10.1002/14651858.CD009618
	Advance care planning for end-stage kidney disease	10.1002/14651858.CD010687
National Guideline Alliance (hosted by the Royal College of Obstetricians and Gynaecologists); National Institute for Health and Care Excellence (commissioner). Pancreatic cancer in adults: diagnosis and management. London: National Institute for Health and Care Excellence; 2018. (NICE NG85). [Issued February 2018]. Available from: https://www.nice.org.uk/guidance/ng85/evidence	Celiac plexus block for pancreatic cancer pain in adults	10.1002/14651858.CD007519.pub2
National Institute for Health and Care Excellence. Clinical Knowledge Summaries: Chronic obstructive pulmonary disease. London: National Institute for Health and Care Excellence; 2018. [last revised December 2018]. Available from: http://cks.nice.org.uk/chronic-obstructive-pulmonary-disease	Benzodiazepines for the relief of breathlessness in advanced malignant and non-malignant diseases in adults	10.1002/14651858.CD007354.pub2
National Institute for Health and Care Excellence. Clinical Knowledge Summaries: Palliative care - oral. London: National Institute for Health and Care Excellence; 2018. [last revised October 2018]. Available from: http://cks.nice.org.uk/palliative-care-oral	Parasympathomimetic drugs for the treatment of salivary gland dysfunction due to radiotherapy	10.1002/14651858.CD003782.pub3
National Institute for Health and Care Excellence. Clinical Knowledge Summaries: Migraine. London: National Institute for Health and Care Excellence; 2018. [last revised February 2018]. Available from: http://cks.nice.org.uk/migraine	Acupuncture for migraine prophylaxis	10.1002/14651858.CD001218.pub2
	Acupuncture for the prevention of episodic migraine	10.1002/14651858.CD001218.pub3
	Sumatriptan (oral route of administration) for acute migraine attacks in adults	10.1002/14651858.CD008615.pub2
	Sumatriptan (intranasal route of administration) for acute migraine attacks in adults	10.1002/14651858.CD009663
	Sumatriptan (rectal route of administration) for acute migraine attacks in adults	10.1002/14651858.CD009664
	Sumatriptan (subcutaneous route of administration) for acute migraine attacks in adults	10.1002/14651858.CD009665
National Institute for Health and Care Excellence. Clinical Knowledge Summaries: Osteoarthritis. London: National Institute for Health and Care Excellence; 2018. [last revised June 2018]. Available from: https://cks.nice.org.uk/osteoarthritis	Topical NSAIDs for chronic musculoskeletal pain in adults	10.1002/14651858.CD007400.pub2
	Salicylate-containing rubefacients for acute and chronic musculoskeletal pain in adults	10.1002/14651858.CD007403.pub3

<p>Mesothelioma Guideline Development Group; British Thoracic Society. BTS Guideline for the investigation and management of malignant pleural mesothelioma. Thorax. 2018 March;73 Suppl1:i1-i30. doi: 10.1136/thoraxjnl-2017-211321. Available from: https://www.brit-thoracic.org.uk/document-library/clinical-information/mesothelioma/mesothelioma-guideline-2018/bts-guideline-for-the-investigation-and-management-of-malignant-pleural-mesothelioma/</p>	<p>Megestrol acetate for treatment of anorexia-cachexia syndrome</p>	<p>10.1002/14651858.CD004310.pub3</p>
	<p>Non-pharmacological interventions for breathlessness in advanced stages of malignant and non-malignant diseases</p>	<p>10.1002/14651858.CD005623.pub2</p>
	<p>Exercise for the management of cancer-related fatigue in adults</p>	<p>10.1002/14651858.CD006145.pub3</p>
<p>Millington GWM, Collins A, Lovell CR, Leslie TA, Yong ASW, Morgan JD, Ajithkumar T, Andrews MJ, Rushbook SM, Coelho RR, Catten SJ, Lee KYC, Skellett AM, Affleck AG, Exton LS, Mohd Mustapa MF, Levell NJ. British Association of Dermatologists' guidelines for the investigation and management of generalized pruritus in adults without an underlying dermatosis, 2018. British Journal of Dermatology. 2018 January;178:34-60. doi: 10.1111/bjd.16117. Available from: http://www.bad.org.uk/shared/get-file.ashx?id=5927&itemtype=document</p>	<p>Pharmacological interventions for pruritus in adult palliative care patients</p>	<p>10.1002/14651858.CD008320</p>
<p>Chartered Society of Physiotherapy. Physiotherapy works: Chronic pain. London: Chartered Society of Physiotherapy; 2018. [Issued February 2018]. Available from: http://www.csp.org.uk/publications/physiotherapy-works-chronic-pain</p>	<p>Physical activity and exercise for chronic pain in adults: an overview of Cochrane Reviews</p>	<p>10.1002/14651858.CD011279.pub3</p>
<p>Goebel A, Atkins RM, Barker C, Cameron H, Cohen H, Cowell F, Eldridge P, Fullilove S, Gillespie S, Guly H, Henderson J, Ikkos G, Jepson F, Johnson M, Lane N, Lewis J, Lewis S, Mani-Babu S, McCabe CS, Parkinson M, Phillip RD, Poole H, Rodham K, Serpell M, Shenker N, Taylor M, Turner-Stokes L, Vivian C, Wilkinson P, Yates B. Complex regional pain syndrome in adults: UK guidelines for diagnosis, referral and management in primary and secondary care. London: Royal College of Physicians; 2018. Available from: https://www.rcplondon.ac.uk/guidelines-policy/complex-regional-pain-syndrome-adults</p>	<p>Psychological therapies for the management of chronic pain (excluding headache) in adults</p>	<p>10.1002/14651858.CD007407.pub2</p>

<p>The Royal College of Surgeons of England; The British Society for Disability and Oral Health. The Oral Management of Oncology Patients Requiring Radiotherapy, Chemotherapy and / or Bone Marrow Transplantation: Clinical Guidelines (updated 2018). London: Faculty of Dental Surgery, Royal College of Surgeons; 2018. Available from: https://www.rcseng.ac.uk/dental-faculties/fds/publications-guidelines/clinical-guidelines/</p>	<p>Parasympathomimetic drugs for the treatment of salivary gland dysfunction due to radiotherapy</p>	<p>10.1002/14651858.CD003782.pub3</p>
<p>National Osteoporosis Society. Vitamin D and Bone Health: A practical clinical guideline for patient management. Bath: National Osteoporosis Society; 2018 December. Available from: https://nos.org.uk/media/100231/nos_vitamin_d_and_bone_health_in_adults_web.pdf</p>	<p>Vitamin D for the treatment of chronic painful conditions in adults</p>	<p>10.1002/14651858.CD007771.pub3</p>
<p>Short Life Working Group for Paediatric Pain. Management of Chronic Pain in Children and Young People: A National Clinical Guideline. Edinburgh: Scottish Government; 2018 March. Available from: https://www.gov.scot/Resource/0053/00533194.pdf</p>	<p>Psychological therapies for the management of chronic and recurrent pain in children and adolescents</p>	<p>10.1002/14651858.CD003968.pub3</p>
	<p>Psychological therapies for the management of chronic and recurrent pain in children and adolescents</p>	<p>10.1002/14651858.CD003968.pub4</p>
	<p>Local anesthetic sympathetic blockade for complex regional pain syndrome</p>	<p>10.1002/14651858.CD004598.pub2</p>
	<p>Local anaesthetic sympathetic blockade for complex regional pain syndrome</p>	<p>10.1002/14651858.CD004598.pub3</p>
	<p>Opioid switching to improve pain relief and drug tolerability</p>	<p>10.1002/14651858.CD004847</p>
	<p>Pharmacological interventions for pain in children and adolescents with life-limiting conditions</p>	<p>10.1002/14651858.CD010750.pub2</p>
	<p>Antidepressants for chronic non-cancer pain in children and adolescents</p>	<p>10.1002/14651858.CD012535.pub2</p>
	<p>Antiepileptic drugs for chronic non-cancer pain in children and adolescents</p>	<p>10.1002/14651858.CD012536.pub2</p>
	<p>Non-steroidal anti-inflammatory drugs (NSAIDs) for chronic non-cancer pain in children and adolescents</p>	<p>10.1002/14651858.CD012537.pub2</p>
	<p>Opioids for chronic non-cancer pain in children and adolescents</p>	<p>10.1002/14651858.CD012538.pub2</p>
	<p>Paracetamol (acetaminophen) for chronic non-cancer pain in children and adolescents</p>	<p>10.1002/14651858.CD012539.pub2</p>

<p>Patel MX, Sethi FN, Barnes TR, Dix R, Dratcu L, Fox B, Garriga M, Haste JC, Kahl KG, Lingford-Hughes A, McAllister-Williams H, O'Brien A, Parker C, Paterson B, Paton C, Posporelis S, Taylor DM, Vieta E, Völlm B, Wilson-Jones C, Woods L. Joint BAP NAPICU evidence-based consensus guidelines for the clinical management of acute disturbance: De-escalation and rapid tranquillisation. J Psychopharmacol. 2018 Jun;32(6):601-640. doi: 10.1177/0269881118776738. Available from: https://www.bap.org.uk/pdfs/BAP_Guidelines-</p>	<p>Levomepromazine for nausea and vomiting in palliative care</p>	<p>10.1002/14651858.CD009420.pub3</p>
<p>Cox S, Dickson J, Goddard A, Hughes J, Land S, Miller B, Rice A; Royal College of Physicians (RCP); Royal College of Radiologists (RCR); Royal College of Radiologists (RCR). Recommendations on cannabis-based products for medicinal use. London: Royal College of Physicians; 2018 October. Available from: https://www.rcplondon.ac.uk/projects/outputs/recommendations-cannabis-based-products-medicinal-use</p>	<p>Cannabis-based medicines for chronic neuropathic pain in adults</p>	<p>10.1002/14651858.CD012182.pub2</p>
<p>Office of Nursing and Midwifery Services Director Quality Care-Metrics Project Group. National Guideline for Nursing and Midwifery Quality Care-Metrics Data Measurement in Public Health Nursing Services 2018. Dublin: Health Service Executive; 2018 December. Available from: https://www.hse.ie/eng/services/publications/nursingmidwifery%20services/national-guideline-qcm-public-health.pdf</p>	<p>Opioids for cancer pain - an overview of Cochrane reviews</p>	<p>10.1002/14651858.CD012592.pub2</p>
<p>Office of Nursing and Midwifery Services Director Quality Care-Metrics Project Group. National Guideline for Nursing and Midwifery Quality Care-Metrics Data Measurement in Acute Care 2018. Dublin: Health Service Executive; 2018 December. Available from: https://www.hse.ie/eng/services/publications/nursingmidwifery%20services/national-guideline-qcm-acute.pdf</p>	<p>Opioids for cancer pain - an overview of Cochrane reviews</p>	<p>10.1002/14651858.CD012592.pub2</p>

<p>Lung Cancer Disease Site Group. A Quality Initiative of the Program in Evidence-Based Care (PEBC), Cancer Care Ontario (CCO): The Role of High Dose Rate Brachytherapy in the Palliation of Symptoms in Patients with Non-Small Cell Lung Cancer (Evidence-based Series 7-16 Version 3). Toronto: Cancer Care Ontario; 2018 June. Available from: https://www.cancercareontario.ca/en/guidelines-advice/types-of-cancer/791</p>	<p>Interventions for cough in cancer</p>	<p>10.1002/14651858.CD007881.pub3</p>
<p>Bussi�eres AE, Stewart G, Al-Zoubi F, Decina P, Descarreaux M, Haskett D, Hincapi�e C, Pag�e I, Passmore S, Srbely J, Stupar M, Weisberg J, Ornelas J. Spinal Manipulative Therapy and Other Conservative Treatments for Low Back Pain: A Guideline From the Canadian Chiropractic Guideline Initiative. J Manipulative Physiol Ther. 2018 May;41(4):265-293. doi: 10.1016/j.jmpt.2017.12.004. Available from: https://www.jmptonline.org/article/S0161-4754(17)30236-1/references</p>	<p>Physical activity and exercise for chronic pain in adults: an overview of Cochrane Reviews</p>	<p>10.1002/14651858.CD011279.pub3</p>
<p>Alberta Provincial Palliative Care Tumour Team, Alberta Health Services. Cancer Pain (Clinical Practice Guideline Supp-012) Version 1. Edmonton: Alberta Health Services; 2018. [Issued September 2018]. Available from: https://www.albertahealthservices.ca/assets/info/hp/cancer/if-hp-cancer-guide-supp-012-cancer-pain.pdf</p>	<p>Ketamine as an adjuvant to opioids for cancer pain</p>	<p>10.1002/14651858.CD003351.pub3</p>
	<p>Corticosteroids for the management of cancer-related pain in adults</p>	<p>10.1002/14651858.CD010756.pub2</p>
<p>Assayag D, Camp PG, Fisher J, Johannson KA, Kolb M, Lohmann T, Manganas H, Morisset J, Ryerson CJ, Shapera S, Simon J, Singer LG, Fell CD; Canadian Thoracic Society. Comprehensive management of fibrotic interstitial lung diseases: A Canadian Thoracic Society position statement. Canadian Journal of Respiratory, Critical Care, and Sleep Medicine. 2018 Sept; 2(4):234-243. doi: 10.1080/24745332.2018.1503456. Available from: https://cts-sct.ca/wp-content/uploads/2019/03/CTS-Comprehensive-management-of-Fibrotic-ILD.pdf</p>	<p>Benzodiazepines for the relief of breathlessness in advanced malignant and non-malignant diseases in adults</p>	<p>10.1002/14651858.CD007354.pub2</p>
<p>Global Initiative for Chronic Obstructive Lung Disease. Global strategy for the diagnosis, management and prevention of chronic obstructive pulmonary disease: 2018 report. Vancouver (WA): Global Initiative for Chronic Obstructive Lung Disease (GOLD);</p>	<p>Non-pharmacological interventions for breathlessness in advanced stages of malignant and non-malignant diseases</p>	<p>10.1002/14651858.CD005623.pub2</p>
	<p>Benzodiazepines for the relief of breathlessness in advanced malignant and non-malignant diseases in adults</p>	<p>10.1002/14651858.CD007354.pub2</p>

<p>2018. Available from: https://goldcopd.org/wp-content/uploads/2017/11/GOLD-2018-v6.0-FINAL-revised-20-</p>	<p>Interventions for fatigue and weight loss in adults with advanced progressive illness</p>	<p>10.1002/14651858.CD008427.pub2</p>
<p>Memtsoudis SG, Cozowicz C, Nagappa M, Wong J, Joshi GP, Wong DT, Doufas AG, Yilmaz M, Stein MH, Krajewski ML, Singh M, Pichler L, Ramachandran SK, Chung F; Society of Anesthesia and Sleep Medicine. Society of Anesthesia and Sleep Medicine Guideline on Intraoperative Management of Adult Patients With Obstructive Sleep Apnea. <i>Anesth Analg</i>. 2018 Oct;127(4):967-987. doi: 10.1213/ANE.0000000000003434. Available from: https://www.ncbi.nlm.nih.gov/pubmed/29944522</p>	<p>Perioperative ketamine for acute postoperative pain</p>	<p>10.1002/14651858.CD004603.pub2</p>
<p>Langman LJ, Jannetto; The American Academy of Pain Medicine. Laboratory Medicine Practice Guidelines: Using Clinical Laboratory Tests to Monitor Drug Therapy in Pain Management Patients. Washington, DC: American Association for Clinical Chemistry; 2018. Available from: https://www.aacc.org/-/media/Files/Science-and-Practice/Practice-Guidelines/Pain-Management/LMPGPain-Management20171220.pdf?la=en&hash=19670524407619F78999AB60731A24CB4901939D</p>	<p>Long-term opioid management for chronic noncancer pain</p>	<p>10.1002/14651858.CD006605.pub2</p>
<p>American College of Radiology. ACR Appropriateness Criteria: Stage I Breast Cancer: Management of Vertebral Compression Fractures. Virginia: American College of Radiology; 2018. Available from: https://acsearch.acr.org/docs/70545/Narrative/</p>	<p>Interventions for the treatment of metastatic extradural spinal cord compression in adults</p>	<p>10.1002/14651858.CD006716.pub3</p>
<p>Cohen SP, Bhatia A, Buvanendran A, Schwenk ES, Wasan AD, Hurley RW, Viscusi ER, Narouze S, Davis FN, Ritchie EC, Lubenow TR, Hooten WM. Consensus Guidelines on the Use of Intravenous Ketamine Infusions for Chronic Pain From the American Society of Regional Anesthesia and Pain Medicine, the American Academy of Pain Medicine, and the American Society of Anesthesiologists. <i>Reg Anesth Pain Med</i>. 2018 Jul;43(5):521-546. doi: 10.1097/AAP.0000000000000808. Available from: https://journals.lww.com/rapm/Fulltext/2018/07000/Consensus_Guidelines_on_the_Use_of_Intravenous.11.aspx</p>	<p>Interventions for treating pain and disability in adults with complex regional pain syndrome</p>	<p>10.1002/14651858.CD009416.pub2</p>

<p>Carter J, Lacchetti C, Andersen B, Barton D, Bolte S, Damast S, Diefenbach M, DuHamel K, Florendo J, Ganz P, Goldfarb S, Hallmeyer S, Kushner D, Rowland J; American Society of Clinical Oncology. Interventions to Address Sexual Problems in People With Cancer: American Society of Clinical Oncology Clinical Practice Guideline Adaptation of Cancer Care Ontario Guideline. <i>Journal of Clinical Oncology</i>. 2018 February;36(5):492-511. doi: 10.1200/JCO.2017.75.8995. Available from: https://www.jco.org/journal/36/5/492</p>	<p>Interventions for sexual dysfunction following treatments for cancer in women</p>	<p>10.1002/14651858.CD005540.pub3</p>
<p>Sparks A, Cohen A, Adjao S, Arnold B, Burdick C, Dumas SEB, Graham J, Greene J, Mayfield R, Purpura P, Rochelle R, Scearce T, Stedronsky A, Sucato G, Ying P. Migraine and Tension Headache Guideline. Seattle (WA): Kaiser Permanente; 2018 April. Available from: https://wa.kaiserpermanente.org/static/pdf/public/guidelines/headache.pdf</p>	<p>Acupuncture for the prevention of tension-type headache</p>	<p>10.1002/14651858.CD007587.pub2</p>
	<p>Selective serotonin reuptake inhibitors (SSRIs) and serotonin-norepinephrine reuptake inhibitors (SNRIs) for the prevention of tension-type headache in adults</p>	<p>10.1002/14651858.CD011681</p>
	<p>Ketoprofen for episodic tension-type headache in adults</p>	<p>10.1002/14651858.CD012190.pub2</p>
<p>Feller-Kopman DJ, Reddy CB, DeCamp MM, Diekemper RL, Gould MK, Henry T, Iyer NP, Lee YCG, Lewis SZ, Maskell NA, Rahman NM, Sterman DH, Wahidi MM, Balekian AA. Management of Malignant Pleural Effusions. An Official ATS/STS/STR Clinical Practice Guideline. <i>Am J Respir Crit Care Med</i>. 2018 Oct;198(7):839-849. doi: 10.1164/rccm.201807-1415ST. Available from: https://www.atsjournals.org/doi/ref/10.1164/rccm.201807-1415ST</p>	<p>Pleurodesis for malignant pleural effusions</p>	<p>10.1002/14651858.CD002916.pub2</p>
	<p>Interventions for the management of malignant pleural effusions: a network meta-analysis</p>	<p>10.1002/14651858.CD010529.pub2</p>
<p>Children's Oncology Group. Long-Term Follow-Up Guidelines for survivors of Childhood, Adolescent and Young Adults Cancers, Version 5.0. Monrovia, CA: Children's Oncology Group; 2018 October. Available from: http://www.survivorshipguidelines.org/pdf/2018/COG_LTFU_Guidelines_v5.pdf</p>	<p>Interventions for sexual dysfunction following treatments for cancer in women</p>	<p>10.1002/14651858.CD005540.pub3</p>
<p>Osman H, Shrestha S, Temin S, Ali ZV, Corvera RA, Ddungu HD, De Lima L, Del Pilar Estevez-Diz M, Ferris FD, Gafer N, Gupta H, Horton S, Jacob G, Jia R, Lu FL, Mosoiu D, Puchalski C, Seigel C, Soyannwo O, Cleary J; American Society of Clinical Oncology. Palliative Care in the Global Setting: ASCO Resource-Stratified Practice Guideline. <i>J Glob Oncol</i>. 2018 Jul;(4):1-24. doi: 10.1200/JGO.18.00026. Available from: https://www.jco.org/journal/36/7/1</p>	<p>Oral morphine for cancer pain</p>	<p>10.1002/14651858.CD003868</p>
	<p>Oral morphine for cancer pain</p>	<p>10.1002/14651858.CD003868.pub2</p>
	<p>Oral morphine for cancer pain</p>	<p>10.1002/14651858.CD003868.pub3</p>
	<p>Oral morphine for cancer pain</p>	<p>10.1002/14651858.CD003868.pub4</p>

National Consensus Project for Quality Palliative Care. Clinical Practice Guidelines for Quality Palliative Care, 4th edition. Richmond, VA: National Coalition for Hospice and Palliative Care; 2018. Available from: https://www.nationalcoalitionhpc.org/wp-content/uploads/2018/10/NCHPC-NCPGuidelines_4thED_web_FINAL.pdf	Haloperidol for the treatment of nausea and vomiting in palliative care patients	10.1002/14651858.CD006271.pub2
Chaikof EL, Dalman RL, Eskandari MK, Jackson BM, Lee WA, Mansour MA, Mastracci TM, Mell M, Murad MH, Nguyen LL, Oderich GS, Patel MS, Schermerhorn ML, Starnes BW. The Society for Vascular Surgery practice guidelines on the care of patients with an abdominal aortic aneurysm. J Vasc Surg. 2018 Jan;67(1):2-77.e2. doi: 10.1016/j.jvs.2017.10.044. Available from: https://www.jvascsurg.org/article/S0741-5214(17)32369-8/references	Perioperative transversus abdominis plane (TAP) blocks for analgesia after abdominal surgery	10.1002/14651858.CD007705.pub2
Yang IA, Dabscheck E, George J, Jenkins S, McDonald CF, McDonald V, Smith B, Zwar N; on behalf of the Lung Foundation Australia and the Thoracic Society of Australia and New Zealand. The COPD-X Plan: Australian and New Zealand Guidelines for the management of Chronic Obstructive Pulmonary Disease 2018. Version 2.56, December 2018. Sydney, NSW, Australia: Lung Foundation Australia and Thoracic Society of Australia and New Zealand; 2018. Available from: http://condx.org.au/	Benzodiazepines for the relief of breathlessness in advanced malignant and non-malignant diseases in adults	10.1002/14651858.CD007354.pub3
	Opioids for the palliation of refractory breathlessness in adults with advanced disease and terminal illness	10.1002/14651858.CD011008.pub2
Department of Health, Government of South Australia. South Australian Perinatal Practice Guidelines: post dural puncture headache. Adelaide, Australia: Department of Health, Government of South Australia; 2018 November. Available from: https://www.sahealth.sa.gov.au/wps/wcm/connect/de1a0e01-5d4a-40ec-97b7-db0816fa8a31/Post+Dural+Puncture+Headache_PPG_v1_0.pdf?MOD=AJPERES&CACHEID=ROOTWORKSPACE-de1a0e01-5d4a-40ec-97b7-db0816fa8a31-mKONh0R	Epidural blood patching for preventing and treating post-dural puncture headache	10.1002/14651858.CD001791.pub2
Deutsche Gesellschaft für Psychosomatische Medizin und Ärztliche Psychotherapie e.V. (DGPM); Deutsches Kollegium für Psychosomatische Medizin (DKPM). S3-Leitlinie: Funktionelle Körperbeschwerden. [S3-Guideline: Functional Body Discomfort]. Berlin: Deutsche Gesellschaft für Psychosomatische Medizin und	Psychological therapies for the management of chronic pain (excluding headache) in adults	10.1002/14651858.CD007407.pub2
	Botulinum toxin for myofascial pain syndromes in adults	10.1002/14651858.CD007533.pub3
	Acupuncture for the prevention of tension-type headache	10.1002/14651858.CD007587.pub2
	Milnacipran for pain in fibromyalgia in adults	10.1002/14651858.CD008244.pub3

<p>Ärztliche Psychotherapie e.V. (DGPM); 2018 July. Available from: https://www.awmf.org/leitlinien/detail/II/051-001.html.</p>	Ibuprofen for acute treatment of episodic tension-type headache in adults	10.1002/14651858.CD011474.pub2
	Selective serotonin reuptake inhibitors (SSRIs) and serotonin-norepinephrine reuptake inhibitors (SNRIs) for the prevention of tension-type headache in adults	10.1002/14651858.CD011681
	Cannabinoids for fibromyalgia	10.1002/14651858.CD011694.pub2
	Pregabalin for pain in fibromyalgia in adults	10.1002/14651858.CD011790.pub2
	Antipsychotics for fibromyalgia in adults	10.1002/14651858.CD011804.pub2
	Aspirin for acute treatment of episodic tension-type headache in adults	10.1002/14651858.CD011888.pub2
	Paracetamol (acetaminophen) for acute treatment of episodic tension-type headache in adults	10.1002/14651858.CD011889.pub2
	Ketoprofen for episodic tension-type headache in adults	10.1002/14651858.CD012190.pub2
	Oral nonsteroidal anti-inflammatory drugs for fibromyalgia in adults	10.1002/14651858.CD012332.pub2
	Milnacipran for neuropathic pain in adults	10.1002/14651858.CD011789
<p>Schulz M, Martin E, Dalhoff K, Worth H, Criée C-P, Lommatzsch M, Hamelmann E, Schneider A, Freitag M, Nowak D, Kraus T, Kainer F, Beule G, Hosemann W, Buhl R, Virchow J-C, Schuster A, Kopp M, Virchow J-C, Hering T, Schultz K, Deter H-C, Pfeiffer-Kascha D, Tholen GR, Vogelberg C, Spindler T, Gappa M, Gerstlauer M, Langhorst J, Klose P, Köhler M, Alsdorf E. Nationale VersorgungsLeithlinie Asthma. [National Care Guideline: Asthma]. Berlin: Ärztliches Zentrum für Qualität in der Medizin (ÄZQ); 2018. Available from: https://www.awmf.org/uploads/tx_szleitlinien/nvl-002l_S3_Asthma_2018-09.pdf</p>	Psychological interventions for parents of children and adolescents with chronic illness	10.1002/14651858.CD009660.pub3
<p>Diener HC, Gaul C, Kropp P; Deutsche Gesellschaft für Neurologie (DGN); Deutsche Migräne- und Kopfschmerzgesellschaft (DMKG). S1 Leitlinien: Therapie der Migräneattacke und Prophylaxe der Migräne. [S1 Guideline: Treatment of migraine attacks and prophylaxis of migraine]. Berlin: Deutsche Gesellschaft für Neurologie (DGN); 2018 January. Available from: https://www.awmf.org/uploads/tx_szleitlinien/030-057l_S1_Migraene-Therapie_2018-04_1.pdf</p>	Acupuncture for migraine prophylaxis	10.1002/14651858.CD001218.pub2
	Acupuncture for the prevention of episodic migraine	10.1002/14651858.CD001218.pub3
	Drugs for preventing migraine headaches in children	10.1002/14651858.CD002761
	Propranolol for migraine prophylaxis	10.1002/14651858.CD003225.pub2
	Dipyrone for acute primary headaches	10.1002/14651858.CD004842.pub2
	Ibuprofen with or without an antiemetic for acute migraine headaches in adults	10.1002/14651858.CD008039.pub3
	Paracetamol (acetaminophen) with or without an antiemetic for acute migraine headaches in adults	10.1002/14651858.CD008040.pub3

Aspirin with or without an antiemetic for acute migraine headaches in adults	10.1002/14651858.CD008041.pub2
Aspirin with or without an antiemetic for acute migraine headaches in adults	10.1002/14651858.CD008041.pub3
Sumatriptan plus naproxen for the treatment of acute migraine attacks in adults	10.1002/14651858.CD008541.pub3
Sumatriptan (oral route of administration) for acute migraine attacks in adults	10.1002/14651858.CD008615.pub2
Zolmitriptan for acute migraine attacks in adults	10.1002/14651858.CD008616.pub2
Sumatriptan (intranasal route of administration) for acute migraine attacks in adults	10.1002/14651858.CD009663
Psychological therapies (Internet-delivered) for the management of chronic pain in adults	10.1002/14651858.CD010152.pub2
Non-prescription (OTC) oral analgesics for acute pain - an overview of Cochrane reviews	10.1002/14651858.CD010794.pub2
Psychological therapies (remotely delivered) for the management of chronic and recurrent pain in children and adolescents	10.1002/14651858.CD011118.pub2
Sumatriptan (rectal route of administration) for acute migraine attacks in adults	10.1002/14651858.CD009664
Sumatriptan (subcutaneous route of administration) for acute migraine attacks in adults	10.1002/14651858.CD009665
Opioids for the palliation of breathlessness in terminal illness	10.1002/14651858.CD002066
Acupuncture-point stimulation for chemotherapy-induced nausea or vomiting	10.1002/14651858.CD002285.pub2
Pleurodesis for malignant pleural effusions	10.1002/14651858.CD002916.pub2
Laxatives or methylnaltrexone for the management of constipation in palliative care patients	10.1002/14651858.CD003448.pub3
Opioids for the management of breakthrough pain in cancer patients	10.1002/14651858.CD004311.pub3
Oxygen therapy for dyspnoea in adults	10.1002/14651858.CD004769.pub2
Selenium for alleviating the side effects of chemotherapy, radiotherapy and surgery in cancer patients	10.1002/14651858.CD005037.pub2
Interventions for noisy breathing in patients near to death	10.1002/14651858.CD005177.pub2

Deutsche Gesellschaft für Pneumologie und Beatmungsmedizin (DGP); Deutsche Krebsgesellschaft (DKG). S3-Leitlinie Prävention, Diagnostik, Therapie und Nachsorge des Lungenkarzinoms. [S3 guideline Prevention, diagnostics, Therapy and follow-up of lung cancer]. Berlin: Leitlinienprogramm Onkologie; 2018 February. Available from: <http://www.awmf.org/leitlinien/detail/II/020-007OL.html>

	Non-pharmacological interventions for breathlessness in advanced stages of malignant and non-malignant diseases	10.1002/14651858.CD005623.pub2
	Benzodiazepines for the relief of breathlessness in advanced malignant and non-malignant diseases in adults	10.1002/14651858.CD007354.pub2
Engeler D, Baranowski AP, Borovicka J, Cottrell A, Dinis-Oliveira P, Elneil S, Hughes J, Messelink EJ, de C Williams AC; European Association of Urology . Guidelines on Chronic Pelvic Pain. Arnhem, The Netherlands: European Association of Urology. [Updated 2018]. Available from: https://uroweb.org/guideline/chronic-pelvic-pain/	Transcutaneous electrical nerve stimulation (TENS) for chronic pain	10.1002/14651858.CD003222.pub2
	Carbamazepine for acute and chronic pain in adults	10.1002/14651858.CD005451.pub2
	Antidepressants for neuropathic pain	10.1002/14651858.CD005454.pub2
	Long-term opioid management for chronic noncancer pain	10.1002/14651858.CD006605.pub2
	Pregabalin for acute and chronic pain in adults	10.1002/14651858.CD007076.pub2
	Psychological therapies for the management of chronic pain (excluding headache) in adults	10.1002/14651858.CD007407.pub3
	Gabapentin for chronic neuropathic pain and fibromyalgia in adults	10.1002/14651858.CD007938.pub2
Leitlinienprogramm Onkologie (Deutsche Krebsgesellschaft, Deutsche Krebshilfe, AWMF). S3-Leitlinie: Diagnostik, Therapie und Nachsorge der Patientinnen mit Endometriumkarzinom. [S3 guideline: diagnosis, treatment and follow-up of patients with endometrial cancer]. Berlin: Leitlinienprogramm Onkologie; 2018 April. Available from: https://www.awmf.org/leitlinien/detail/II/032-034OL.html .	Exercise for the management of cancer-related fatigue in adults	10.1002/14651858.CD006145.pub3
Fallon M, Giusti R, Aielli F, Hoskin P, Rolke R, Sharma M, Ripamonti CI; ESMO Guidelines Committee. Management of cancer pain in adult patients: ESMO Clinical Practice Guidelines. Ann Oncol. 2018 Oct;29(Supplement_4):iv166-iv191. doi: 10.1093/annonc/mdy152. Available from: https://academic.oup.com/annonc/article/29/Supplement_4/iv166/5046945	Radioisotopes for metastatic bone pain	10.1002/14651858.CD003347.pub2
	Ketamine as an adjuvant to opioids for cancer pain	10.1002/14651858.CD003351.pub3
	Laxatives or methylnaltrexone for the management of constipation in palliative care patients	10.1002/14651858.CD003448.pub3
	Oral morphine for cancer pain	10.1002/14651858.CD003868.pub4
	Oxycodone for cancer-related pain	10.1002/14651858.CD003870.pub5
	Codeine, alone and with paracetamol (acetaminophen), for cancer pain	10.1002/14651858.CD006601.pub4
	Celiac plexus block for pancreatic cancer pain in adults	10.1002/14651858.CD007519.pub2
	Spinal cord stimulation for cancer-related pain in adults	10.1002/14651858.CD009389.pub3
	Transdermal fentanyl for cancer pain	10.1002/14651858.CD010270.pub2
	Tramadol with or without paracetamol (acetaminophen) for cancer pain	10.1002/14651858.CD012508.pub2
	Opioids for cancer pain - an overview of Cochrane reviews	10.1002/14651858.CD012592.pub2
	Oral paracetamol (acetaminophen) for cancer pain	10.1002/14651858.CD012637.pub2

	Oral nonsteroidal anti-inflammatory drugs (NSAIDs) for cancer pain in adults	10.1002/14651858.CD012638.pub2
Stöve J; Deutsche Gesellschaft für Orthopädie und Orthopädische Chirurgie (DGOOC). S2k-Leitlinie: Gonarthrose. [S2K Guideline: Gonarthrosis] .Berlin: Deutsche Gesellschaft für Orthopädie und Orthopädische Chirurgie e. V. (DGOOC); 2018 January. Available from: http://www.awmf.org/leitlinien/detail/II/033-004.html .	Topical NSAIDs for chronic musculoskeletal pain in adults	10.1002/14651858.CD007400.pub2
Glocker F. et al; Deutsche Gesellschaft für Neurologie (DGN). S2k Leitlinie: Lumbale Radikulopathie. [S2k Guideline: Lumbar Radiculopathy]. Berlin: Deutsche Gesellschaft für Neurologie (DGN); 2018 January. Available from: https://www.awmf.org/uploads/tx_szleitlinien/030-059L_S2k_Lumbale_Radikulopathie_2018_04.pdf	Carbamazepine for chronic neuropathic pain and fibromyalgia in adults	10.1002/14651858.CD005451.pub3
	Long-term opioid management for chronic noncancer pain	10.1002/14651858.CD006605.pub2
	Pregabalin for acute and chronic pain in adults	10.1002/14651858.CD007076.pub2
Planchard D, Popat S, Kerr K, Novello S, Smit EF, Faivre-Finn C, Mok TS, Reck M, Van Schil PE, Hellmann MD, Peters S; ESMO Guidelines Committee. Metastatic non-small cell lung cancer: ESMO Clinical Practice Guidelines for diagnosis, treatment and follow-up. Ann Oncol. 2018 Oct;29(Supplement_4):iv192-iv237. doi: 10.1093/annonc/mdy275. Available from: https://academic.oup.com/annonc/article/29/Supplement_4/iv192/5115264	Pleurodesis for malignant pleural effusions	10.1002/14651858.CD002916.pub2
	Interventions for the treatment of metastatic extradural spinal cord compression in adults	10.1002/14651858.CD006716.pub3
Leitlinienprogramm Onkologie (Deutsche Krebsgesellschaft, Deutsche Krebshilfe, AWMF): Interdisziplinäre Leitlinie der Qualität S3 zur Früherkennung, Diagnose und Therapie der verschiedenen Stadien des Prostatakarzinoms. [S3 Interdisciplinary Guideline for early detection, diagnosis and treatment of different stages of prostate cancer]. Berlin: Leitlinienprogramm Onkologie; 2018 April. Available from: https://www.awmf.org/leitlinien/detail/II/043-022OL.html	Interventions for the treatment of metastatic extradural spinal cord compression in adults	10.1002/14651858.CD006716.pub2
Wordliczek J, Kotlinska-Lemieszek A, Leppert W, Woron J, Dobrogowski J, Krajnik M, Przeklasa-Muszynska A, Jassem J, Drobnik J, Wrzosek A, Janecki M, Pyszkowska J, Kocot-Kepska M, Zajackowska R, Filipczak-Bryniarska I, Boczar K, Jakowicka-Wordliczek J, Malec-Milewska M, Kübler A, Suchorzewski M,	Ketamine as an adjuvant to opioids for cancer pain	10.1002/14651858.CD003351.pub2
	Oral morphine for cancer pain	10.1002/14651858.CD003868.pub4
	Oxycodone for cancer-related pain	10.1002/14651858.CD003870.pub5
	Opioids for the management of breakthrough pain in cancer patients	10.1002/14651858.CD004311.pub3

<p>Mordarski S, Dziki A, Pasnik K. Pharmacotherapy of pain in cancer patients - recommendations of the Polish Association for the Study of Pain, Polish Society of Palliative Medicine, Polish Society of Oncology, Polish Society of Family Medicine, Polish Society of Anaesthesiology and Intensive Therapy and Association of Polish Surgeons. Pol Przegl Chir. 2018 Aug;90(4):55-84. doi: 10.5604/01.3001.0012.2904. Available from: https://ppch.pl/resources/html/article/details?id=176785&language=en.</p>	Interventions for the treatment of metastatic extradural spinal cord compression in adults	10.1002/14651858.CD006716.pub3
	Buprenorphine for treating cancer pain	10.1002/14651858.CD009596.pub4
	Transdermal fentanyl for cancer pain	10.1002/14651858.CD010270.pub2
	Corticosteroids for the management of cancer-related pain in adults	10.1002/14651858.CD010756.pub2
	Opioids for the palliation of refractory breathlessness in adults with advanced disease and terminal illness	10.1002/14651858.CD011008.pub2
	Hydromorphone for cancer pain	10.1002/14651858.CD011108.pub2
	Tramadol with or without paracetamol (acetaminophen) for cancer pain	10.1002/14651858.CD012508.pub2
<p>Sundhedsstyrelsen, Danish Health Authority. National klinisk retningslinje: Udredning og behandling samt rehabilitering af patienter med generaliserede smerter i bevægeapparatet. [National Clinical Guideline for the treatment and rehabilitation of patients with generalised pain on movement]. Copenhagen: Sundhedsstyrelsen, Danish Health Authority; June, 2018. Available from: https://www.sst.dk/da/udgivelser/2018/~media/82268C6BAF19461399C4FC8D1647DCC0.ashx</p>	Psychological therapies for the management of chronic pain (excluding headache) in adults	10.1002/14651858.CD007407.pub3
	Amitriptyline for fibromyalgia in adults	10.1002/14651858.CD011824
<p>Jara C, Del Barco S, Grávalos C, Hoyos S, Hernández B, Muñoz M, Quintanar T, Meana JA, Rodríguez C, de Las Peñas R. SEOM clinical guideline for treatment of cancer pain (2017). Clin Transl Oncol. 2018 Jan;20(1):97-107. doi: 10.1007/s12094-017-1791-2. Available from: https://www.ncbi.nlm.nih.gov/pmc/articles/PMC5785609/</p>	Methadone for cancer pain	10.1002/14651858.CD003971.pub4
	Hydromorphone for cancer pain	10.1002/14651858.CD011108.pub2
	Oral nonsteroidal anti-inflammatory drugs (NSAIDs) for cancer pain in adults	10.1002/14651858.CD012638.pub2
<p>Deutschsprachige Medizinische Gesellschaft für Paraplegiologie e.V. (DMGP). S2k Leitlinie: Schmerzen bei Querschnittlähmung. [S2k guideline: Pain in Paraplegia]. Berlin: Deutschsprachige Medizinische Gesellschaft für Paraplegiologie e.V. (DMGP); 2018 May. Available from: https://www.awmf.org/leitlinien/detail/II/179-006.html</p>	Antidepressants for neuropathic pain	10.1002/14651858.CD005454.pub2
	Lamotrigine for acute and chronic pain	10.1002/14651858.CD006044.pub3
	Transcutaneous electrical nerve stimulation for acute pain	10.1002/14651858.CD006142.pub2
	Transcutaneous electrical nerve stimulation (TENS) for phantom pain and stump pain following amputation in adults	10.1002/14651858.CD007264.pub3
	Antiepileptic drugs for neuropathic pain and fibromyalgia - an overview of Cochrane reviews	10.1002/14651858.CD010567.pub2
	Levetiracetam for neuropathic pain in adults	10.1002/14651858.CD010943.pub2
	Physical activity and exercise for chronic pain in adults: an overview of Cochrane Reviews	10.1002/14651858.CD011279.pub3

Leitlinienprogramm Onkologie (Deutsche Krebsgesellschaft, Deutsche Krebshilfe, AWMF). S3 Leitlinie: Diagnostik, Therapie und Nachsorge des Melanoms. Version 3.1. [S3 guideline: diagnosis, treatment and follow-up of melanomas. Version 3.1]. Berlin: Leitlinienprogramm Onkologie; 2018 July. Available from: https://www.awmf.org/leitlinien/detail/II/032-024OL.html	Acupuncture-point stimulation for chemotherapy-induced nausea or vomiting	10.1002/14651858.CD002285.pub2
	Eicosapentaenoic acid (EPA, an omega-3 fatty acid from fish oils) for the treatment of cancer cachexia	10.1002/14651858.CD004597.pub2
	Homeopathic medicines for adverse effects of cancer treatments	10.1002/14651858.CD004845.pub2
HerniaSurge Group. International guidelines for groin hernia management. <i>Hernia</i> . 2018 Feb;22(1):1-165. doi: 10.1007/s10029-017-1668-x. Available from: https://www.ncbi.nlm.nih.gov/pmc/articles/PMC5809582/	Topical capsaicin (high concentration) for chronic neuropathic pain in adults	10.1002/14651858.CD007393.pub3
	Perioperative transversus abdominis plane (TAP) blocks for analgesia after abdominal surgery	10.1002/14651858.CD007705
Deutsche Gesellschaft für Gynäkologie und Geburtshilfe (DGGG); Österreichische Gesellschaft für Gynäkologie und Geburtshilfe (OEGGG); Schweizerische Gesellschaft für Gynäkologie und Geburtshilfe (SGGG). S2k Leitlinie: Diagnostik, Therapie und Nachsorge des Vaginalkarzinoms und seiner Vorstufen. [S2k Guideline: Diagnosis, treatment and follow-up of vaginal carcinoma and its precursors]. Berlin: Deutsche Gesellschaft für Gynäkologie und Geburtshilfe (DGGG); 2018 October. Available from: https://www.awmf.org/uploads/tx_szleitlinien/032-042l_S2k_Vaginalkarzinom-Vorstufen-Diagnostik-Therapie-Nachsorge_2018-11.pdf	Selenium for alleviating the side effects of chemotherapy, radiotherapy and surgery in cancer patients	10.1002/14651858.CD005037.pub2
	Psychosocial interventions for reducing fatigue during cancer treatment in adults	10.1002/14651858.CD006953.pub2
Rennekampff HO; Deutschen Gesellschaft für Verbrennungsmedizin (DGV). S2k Leitlinie: Behandlung thermischer Verletzungen des Erwachsenen. [S2k guideline: treatment of thermal injuries in adults]. Berlin: Deutsche Gesellschaft für Verbrennungsmedizin e.V. (DGV); 2018 August. Available from: https://www.awmf.org/uploads/tx_szleitlinien/044-001l_S2k_Thermische__Verletzungen_Erwachsene_2018-12.pdf	Perioperative ketamine for acute postoperative pain	10.1002/14651858.CD004603.pub2
	Intravenous lidocaine for the treatment of background or procedural burn pain	10.1002/14651858.CD005622.pub4
	Pregabalin for acute and chronic pain in adults	10.1002/14651858.CD007076.pub2

<p>Sundhedsstyrelsen, Danish Health Authority. National klinisk retningslinje: Opioidbehandling af kroniske non-maligne smerter. [National Clinical Guideline on opioid treatment of chronic non-cancer pain]. Copenhagen: Sundhedsstyrelsen, Danish Health Authority; December, 2018. Available from: https://www.sst.dk/da/udgivelser/2018/~media/1DC940E1B0C94614B5C15782D6869975.ashx</p>	<p>Physical activity and exercise for chronic pain in adults: an overview of Cochrane Reviews</p>	<p>10.1002/14651858.CD011279.pub3</p>
<p>Suomalaisen Lääkäriseuran Duodecimin; Suomen Ortopediyhdistys ry:n asettama työryhmä; Finnish Medical Society Duodecim; Finnish Orthopaedic Society. Duodecim Käypä Hoito: Polvi- ja lonkkanivelrikkko. [Duodecim Current Care Guidelines: knee and hip osteoarthritis]. Helsinki: Finnish Medical Society Duodecim; May 2018. Available from: http://www.kaypahoito.fi/web/kh/suosituksset/suositus?id=hoi50054</p>	<p>Topical analgesics for acute and chronic pain in adults - an overview of Cochrane Reviews</p>	<p>10.1002/14651858.CD008609.pub2</p>
<p>Suomalaisen Lääkäriseuran Duodecimin; Suomen Palliativisen Lääketieteen yhdistyksen asettama työryhmä; Finnish Medical Society Duodecim; Finnish Palliative Medicine Association. Duodecim Käypä Hoito: Palliativinen hoito ja saattohoito. [Duodecim Current Care Guidelines: palliative treatment and terminal care]. Helsinki: Finnish Medical Society Duodecim; February 2018. Available from: http://www.terveysportti.fi/xmedia/hoi/hoi50063.pdf</p>	<p>Corticosteroids for the resolution of malignant bowel obstruction in advanced gynaecological and gastrointestinal cancer</p>	<p>10.1002/14651858.CD001219</p>
	<p>Opioids for the palliation of breathlessness in terminal illness</p>	<p>10.1002/14651858.CD002066</p>
	<p>Bisphosphonates for the relief of pain secondary to bone metastases</p>	<p>10.1002/14651858.CD002068</p>
	<p>Pleurodesis for malignant pleural effusions</p>	<p>10.1002/14651858.CD002916.pub2</p>
	<p>Calcitonin for metastatic bone pain</p>	<p>10.1002/14651858.CD003223.pub2</p>
	<p>Radioisotopes for metastatic bone pain</p>	<p>10.1002/14651858.CD003347.pub3</p>
	<p>Ketamine as an adjuvant to opioids for cancer pain</p>	<p>10.1002/14651858.CD003351.pub3</p>
	<p>Laxatives or methylalnaltrexone for the management of constipation in palliative care patients</p>	<p>10.1002/14651858.CD003448.pub3</p>
	<p>Laxatives for the management of constipation in people receiving palliative care</p>	<p>10.1002/14651858.CD003448.pub4</p>
	<p>Oral morphine for cancer pain</p>	<p>10.1002/14651858.CD003868.pub4</p>
	<p>Oxycodone for cancer-related pain</p>	<p>10.1002/14651858.CD003870.pub5</p>
<p>Methadone for cancer pain</p>	<p>10.1002/14651858.CD003971.pub3</p>	
<p>Megestrol acetate for the treatment of anorexia-cachexia syndrome</p>	<p>10.1002/14651858.CD004310.pub2</p>	

Megestrol acetate for treatment of anorexia-cachexia syndrome	10.1002/14651858.CD004310.pub3
Opioids for the management of breakthrough (episodic) pain in cancer patients	10.1002/14651858.CD004311.pub2
Opioids for the management of breakthrough pain in cancer patients	10.1002/14651858.CD004311.pub3
Drug therapy for symptoms associated with anxiety in adult palliative care patients	10.1002/14651858.CD004596.pub2
Oxygen therapy for dyspnoea in adults	10.1002/14651858.CD004769.pub2
Drug therapy for delirium in terminally ill adult patients	10.1002/14651858.CD004770.pub2
Interventions for noisy breathing in patients near to death	10.1002/14651858.CD005177.pub2
Antidepressants for neuropathic pain	10.1002/14651858.CD005454.pub2
Psychotherapy for depression among incurable cancer patients	10.1002/14651858.CD005537.pub2
Non-pharmacological interventions for breathlessness in advanced stages of malignant and non-malignant diseases	10.1002/14651858.CD005623.pub2
Haloperidol for the treatment of nausea and vomiting in palliative care patients	10.1002/14651858.CD006271.pub2
Medically assisted hydration for adult palliative care patients	10.1002/14651858.CD006273.pub3
Medically assisted nutrition for palliative care in adult patients	10.1002/14651858.CD006274.pub2
Pharmacologic interventions for treating phantom limb pain	10.1002/14651858.CD006380.pub3
Codeine, alone and with paracetamol (acetaminophen), for cancer pain	10.1002/14651858.CD006601.pub4
Pharmacological treatments for fatigue associated with palliative care	10.1002/14651858.CD006788.pub3
Benzodiazepines for the relief of breathlessness in advanced malignant and non-malignant diseases in adults	10.1002/14651858.CD007354.pub3
Topical capsaicin (high concentration) for chronic neuropathic pain in adults	10.1002/14651858.CD007393.pub4
Gabapentin for chronic neuropathic pain in adults	10.1002/14651858.CD007938.pub4
Combination pharmacotherapy for the treatment of neuropathic pain in adults	10.1002/14651858.CD008943.pub2

	Blood transfusions for anaemia in patients with advanced cancer	10.1002/14651858.CD009007.pub2
	Spinal cord stimulation for cancer-related pain in adults	10.1002/14651858.CD009389.pub3
	Palliative pharmacological sedation for terminally ill adults	10.1002/14651858.CD010206.pub2
	Antiepileptic drugs for neuropathic pain and fibromyalgia - an overview of Cochrane reviews	10.1002/14651858.CD010567.pub2
	Opioids for the palliation of refractory breathlessness in adults with advanced disease and terminal illness	10.1002/14651858.CD011008.pub2
	Venlafaxine for neuropathic pain in adults	10.1002/14651858.CD011091.pub2
	Tramadol with or without paracetamol (acetaminophen) for cancer pain	10.1002/14651858.CD012508.pub2
	Oral paracetamol (acetaminophen) for cancer pain	10.1002/14651858.CD012637.pub2
	Oral nonsteroidal anti-inflammatory drugs (NSAIDs) for cancer pain in adults	10.1002/14651858.CD012638.pub2
NHG Werkgroep Pijn. NHG-Standaard Pijn (Standaard M106). [Dutch College of General Practitioners - Guideline on pain (Guideline M106)]. Utrecht: Nederlands Huisartsen Genootschap; 2018. Available from: https://www.nhg.org/standaarden/volledig/nhg-standaard-pijn	Methadone for cancer pain	10.1002/14651858.CD003971.pub3
	Opioids for the management of breakthrough pain in cancer patients	10.1002/14651858.CD004311.pub3
	Opioid switching to improve pain relief and drug tolerability	10.1002/14651858.CD004847
	Carbamazepine for chronic neuropathic pain and fibromyalgia in adults	10.1002/14651858.CD005451.pub3
	Diclofenac for acute pain in children	10.1002/14651858.CD005538.pub2
	Opioids for neuropathic pain	10.1002/14651858.CD006146.pub2
	Long-term opioid management for chronic noncancer pain	10.1002/14651858.CD006605.pub2
	Pregabalin for acute and chronic pain in adults	10.1002/14651858.CD007076.pub2
	Topical capsaicin (high concentration) for chronic neuropathic pain in adults	10.1002/14651858.CD007393.pub3
	Topical NSAIDs for chronic musculoskeletal pain in adults	10.1002/14651858.CD007400.pub2
	Topical NSAIDs for acute pain in adults	10.1002/14651858.CD007402.pub2
	Psychological therapies for the management of chronic pain (excluding headache) in adults	10.1002/14651858.CD007407.pub3
	Gabapentin for chronic neuropathic pain and fibromyalgia in adults	10.1002/14651858.CD007938.pub3
	Methadone for chronic non-cancer pain in adults	10.1002/14651858.CD008025.pub2
	Amitriptyline for neuropathic pain and fibromyalgia in adults	10.1002/14651858.CD008242.pub2

	Combination pharmacotherapy for the treatment of neuropathic pain in adults	10.1002/14651858.CD008943.pub2
	Topical capsaicin (low concentration) for chronic neuropathic pain in adults	10.1002/14651858.CD010111
	Single dose oral ibuprofen plus paracetamol (acetaminophen) for acute postoperative pain	10.1002/14651858.CD010210.pub2
	Transdermal fentanyl for cancer pain	10.1002/14651858.CD010270.pub2
	Interventions for the reduction of prescribed opioid use in chronic non-cancer pain	10.1002/14651858.CD010323.pub3
	Imipramine for neuropathic pain in adults	10.1002/14651858.CD010769.pub2
	Nortriptyline for neuropathic pain in adults	10.1002/14651858.CD011209.pub2
	Adverse events associated with medium- and long-term use of opioids for chronic non-cancer pain: an overview of Cochrane Reviews	10.1002/14651858.CD012509.pub2
	Opioids for cancer pain - an overview of Cochrane reviews	10.1002/14651858.CD012592.pub2
Hoffmann J, Mollan SP, Paemeleire K, Lampl C, Jensen RH5, Sinclair AJ. European headache federation guideline on idiopathic intracranial hypertension. J Headache Pain. 2018 Oct;19(1):93. doi: 10.1186/s10194-018-0919-2. Available from: https://www.ncbi.nlm.nih.gov/pubmed/30298346	Botulinum toxins for the prevention of migraine in adults	10.1002/14651858.CD011616.pub2
Bush SH; Lawlor PG; Ryan K; Centeno C; Lucchesi M; Kanji S; Siddiqi N; Morandi A; Davis DHJ; Laurent M; Schofield N; Barallat E; Ripamonti CI; ESMO Guidelines Committee. Delirium in adult cancer patients: ESMO Clinical Practice Guidelines. Annals of Oncology. 2018 Oct;29(Supp 4):iv143-iv165. doi: 10.1093/annonc/mdy147. Available from: https://academic.oup.com/annonc/article/29/Supplement_4/iv143/5046944	Palliative pharmacological sedation for terminally ill adults	10.1002/14651858.CD010206.pub2
Larkin PJ, Cherny NI, La Carpia D, Guglielmo M, Ostgathe C, Scotte F, Ripamonti CI; on behalf of the ESMO Guidelines Committee.	Laxatives or methylnaltrexone for the management of constipation in palliative care patients	10.1002/14651858.CD003448.pub3

<p>Diagnosis, Assessment and Management of Constipation in Advanced Cancer: ESMO Clinical Practice Guidelines. <i>Ann Oncol</i>. 2018 Oct;29(Supplement_4):iv111-iv125. doi: 10.1093/annonc/mdy148. Available from: https://www.esmo.org/Guidelines/Supportive-and-Palliative-Care/Diagnosis-Assessment-and-Management-of-Constipation-in-</p>	<p>Laxatives for the management of constipation in people receiving palliative care</p>	<p>10.1002/14651858.CD003448.pub4</p>
<p>Dieterich M, et al.; Deutsche Gesellschaft für Neurologie (DGN). S1 Leitlinien für Diagnostik und Therapie in der Neurologie: postpunktionellen und spontanen Liquorunterdruck-Syndroms. [S1 guideline for diagnosis and treatment in neurology: post-puncture and spontaneous CSF hypertension syndrome]. Berlin: Deutsche Gesellschaft für Neurologie (DGN); 2018 March. Available from: https://www.awmf.org/uploads/tx_szleitlinien/030-113l_S1_Diagnostik-Therapie-postpunktioneller-spontaner-Liquorunterdruck-Syndroms_2018-12.pdf</p>	<p>Epidural blood patching for preventing and treating post-dural puncture headache</p>	<p>10.1002/14651858.CD001791.pub2</p>
	<p>Drug therapy for treating post-dural puncture headache</p>	<p>10.1002/14651858.CD007887.pub2</p>
<p>Song KH, Jung HK, Kim HJ, Koo HS, Kwon Y, Shin HD, Lim HC, Shin JE, Kim SE, Cho DH, Kim JH, Kim HJ; Clinical Practice Guidelines Group Under the Korean Society of Neurogastroenterology and Motility. Clinical Practice Guidelines for Irritable Bowel Syndrome in Korea, 2017 Revised Edition. <i>J Neurogastroenterol Motil</i>. 2018 Apr;24(2):197-215. doi: 10.5056/jnm17145. Available from: https://www.ncbi.nlm.nih.gov/pmc/articles/PMC5885719/.</p>	<p>Antidepressants for neuropathic pain</p>	<p>10.1002/14651858.CD005454.pub2</p>
<p>Lai K, Shen H, Zhou X, Qiu Z, Cai S, Huang K, Wang Q, Wang C, Lin J, Hao C, Kong L, Zhang S, Chen Y, Luo W, Jiang M, Xie J, Zhong N. Clinical Practice Guidelines for Diagnosis and Management of Cough-Chinese Thoracic Society (CTS) Asthma Consortium. <i>J Thorac Dis</i>. 2018 Nov;10(11):6314-6351. doi: 10.21037/jtd.2018.09.153. Available from: https://pubmed.ncbi.nlm.nih.gov/30622806/</p>	<p>Interventions for cough in cancer</p>	<p>10.1002/14651858.CD007881.pub2</p>
<p>Rafanan BS Jr, Valdecañas BF, Lim BP, Malairungsakul A, Tassanawipas W, Shiya C, Tse LF, Luong TK. Consensus</p>	<p>Topical NSAIDs for chronic musculoskeletal pain in adults</p>	<p>10.1002/14651858.CD007400.pub3</p>
	<p>Topical NSAIDs for acute musculoskeletal pain in adults</p>	<p>10.1002/14651858.CD007402.pub3</p>

recommendations for managing osteoarthritic pain with topical NSAIDs in Asia-Pacific. Pain Manag. 2018 Mar;8(2):115-128. doi: 10.2217/pmt-2017-0047. Available from: <https://www.futuremedicine.com/doi/10.2217/pmt-2017-0047>

Salicylate-containing rubefaciants for acute and chronic musculoskeletal pain in adults

10.1002/14651858.CD007403.pub3

Summary of NICE & SIGN Guideline data

74 reviews in: 12 guidelines (9 NICE (1 updated in this period); 3 SIGN) and 7 Clinical Knowledge Summaries (2 updated in this period)

11 reviews in more than one guideline (including updates) (frequency: 1 in 4; 10 in 2)